

Acta Crystallographica Section E

**Structure Reports**

**Online**

ISSN 1600-5368

Editors: **W. Clegg** and **D. G. Watson**

## **Piperazinium dichromate(VI)**

**Bikshandarkoil R. Srinivasan, Ashish R. Naik, Christian Näther and Wolfgang Bensch**

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site provided that this cover page is retained. Republication of this article or its storage in electronic databases or the like is not permitted without prior permission in writing from the IUCr.

Bikshandarkoil R. Srinivasan,<sup>a</sup>  
Ashish R. Naik,<sup>a</sup> Christian  
Näther<sup>b\*</sup> and Wolfgang Bensch<sup>b</sup><sup>a</sup>Department of Chemistry, Goa University  
PO, Goa 403 206, India, and <sup>b</sup>Institut für  
Anorganische Chemie, Christian-Albrechts-  
Universität Kiel, Olshausenstraße 40, D-24098  
Kiel, GermanyCorrespondence e-mail:  
cnaether@ac.uni-kiel.de

## Key indicators


Single-crystal X-ray study  
T = 293 K  
Mean  $\sigma(\text{C}-\text{C}) = 0.003 \text{ \AA}$   
R factor = 0.026  
wR factor = 0.078  
Data-to-parameter ratio = 21.2For details of how these key indicators were  
automatically derived from the article, see  
<http://journals.iucr.org/e>.

## Piperazinium dichromate(VI)

The structure of the title complex,  $(\text{C}_4\text{H}_{12}\text{N}_2)[\text{Cr}_2\text{O}_7]$ , consists of dichromate dianions which are connected to the cyclic organic piperazinium  $(\text{pipH}_2)^{2+}$  dications *via* hydrogen bonding. There are two crystallographically independent piperazinium dications, both located on centres of inversion, while the anions are located in general positions.


## Comment

The reported synthesis and structural characterization of complexes such as  $[\text{M}(\text{py})_4\text{Cr}_2\text{O}_7]$  ( $\text{M} = \text{Cu}, \text{Zn}$ ;  $\text{py} = \text{pyridine}$ ), which exhibit non-linear optical properties (Norquist *et al.*, 2001), is an important reason for the renewed interest in the chemistry of dichromate complexes associated with organic cations. Dichromates are well known for their use in organic syntheses to effect a variety of synthetic transformations (House, 1977). It has been reported that the type of the ammonium cation associated with the dichromate determines the oxidizing property of the dichromate ion and this is inversely related to the donor strength of the associated amine ligand (Santaniello *et al.*, 1983). The clue to the differing oxidizing property lies in understanding the nature of cation–anion interactions. In our work, we have been investigating the synthesis and structural characterization of materials resulting from the interaction of organic diamines with group(VI) oxo and thiometalates. All compounds hitherto reported by us exhibit weak hydrogen-bonding interactions between the organic cation and the anion. We have previously described the structures of several tetrathiomolybdates (Srinivasan *et al.*, 2004, and references therein), tetrathiotungstates (Srinivasan *et al.*, 2003a, and references therein), piperazinium chromate (Srinivasan *et al.*, 2003b) and ethylenediammonium dichromate (Srinivasan *et al.*, 2003c). In continuation of this work, we describe the structure of piperazinium dichromate. In separate research, this complex was shown to be a mild reagent for the oxidation of thiols to disulfides (Movassagh *et al.*, 2002).




The structure of the title compound, (I), consists of  $[\text{Cr}_2\text{O}_7]^{2-}$  dianions and cyclic piperazinium dications (Fig. 1). There are two crystallographically independent cations, both located on centres of inversion, while the anions are located in general positions. As expected, the piperazinium dication

Received 31 August 2004  
Accepted 1 September 2004  
Online 11 September 2004



**Figure 1**

The crystal structure of piperazinium dichromate(VI), with labelling and displacement ellipsoids drawn at the 50% probability level [symmetry codes: (i)  $2 - x, 2 - y, 1 - z$ ; (ii)  $1 - x, 1 - y, -z$ ].



**Figure 2**

The crystal structure of piperazinium dichromate(VI), viewed along the *a* axis (intermolecular hydrogen bonds are shown as dashed lines).

adopts the chair conformation with bond lengths and angles (Table 1) in the range usually observed in this form (Srinivasan *et al.*, 2003*b*; Tran Qui & Palacios, 1990). The O—Cr—O bond angles range from 106.69 (7) to 111.67 (8)° (Table 1), while the terminal Cr—O bond distances in (I) vary from 1.5939 (16) to 1.6334 (14) Å, with a mean terminal Cr—O bond length of 1.6146 Å. The average bridging Cr—O bond distance is 1.7860 Å. These values are in good agreement with those reported for other dichromates (Srinivasan *et al.*, 2003*c*; Panagiotopoulos & Brown, 1972). The terminal Cr—O bond lengths in (I) cover a wider range than in the related complex (enH<sub>2</sub>)[Cr<sub>2</sub>O<sub>7</sub>] (enH<sub>2</sub> is ethylenediammonium), where the

terminal Cr—O bond distances vary from 1.6075 (15) to 1.6236 (14) Å. It is also noted that the difference between the longest and the shortest terminal Cr—O bonds is of the order of 0.0395 Å, whereas in (enH<sub>2</sub>)[Cr<sub>2</sub>O<sub>7</sub>] the difference amounts to 0.0161 Å. Such differences can be attributed to the differing nature of the cations in the two complexes.

The cations and anions are connected *via* hydrogen bonding between the O atoms of the dianion and the H atoms bound to the nitrogen of the cyclic cation (Fig. 2). Short H···O contacts ranging from 1.91 to 2.56 Å are observed in (I) between the O1/2/5/6/7 atoms and the H atoms of the piperazinium cation (Table 2). The shortest H···O separation of 1.91 Å in (I) is slightly shorter than that in (enH<sub>2</sub>)[Cr<sub>2</sub>O<sub>7</sub>], which has a shortest contact of 1.95 Å. Atom O2 in (I) is involved in two hydrogen bonds, which explains the elongation of the Cr1—O2 distance, the longest terminal Cr—O bond in (I). Atoms O6 and O7 each have a single short hydrogen-bonding contact of 1.91 and 1.93 Å, respectively, which are accompanied by large *D—H···A* angles (Table 2). Hence, the observed Cr2—O6 and Cr2—O7 bond lengths of 1.6189 (16) and 1.6242 (14) Å are due to the relatively strong O6···H and O7···H interactions. A long H···O distance of 2.36 Å and a small *D—H···A* angle of 121° indicate a very weak interaction and consequently the Cr2—O5 bond is the shortest [1.5939 (16) Å]. The intermediate Cr1—O1 bond length of 1.6068 (14) Å can be similarly explained based on the strength of the hydrogen-bonding interaction.

## Experimental

(NH<sub>4</sub>)<sub>2</sub>[Cr<sub>2</sub>O<sub>7</sub>] (1.512 g) was dissolved in distilled water (10 ml) containing 0.3 ml of concentrated H<sub>2</sub>SO<sub>4</sub> and this solution was slowly added to a solution of piperazine hexahydrate (1.2 g) dissolved in water (4 ml), resulting in the formation of yellow–orange crystalline blocks of the title complex. The crystals were filtered off, washed with ice-cold water (1 ml) and dried in air (yield 85%). The crystals are quite stable in air. Analysis calculated for C<sub>4</sub>H<sub>12</sub>Cr<sub>2</sub>N<sub>2</sub>O<sub>7</sub>: C 15.79, H 3.98, N 9.21%; found: C 15.17, H 4.11, N 9.39%. <sup>1</sup>H NMR (D<sub>2</sub>O, 300 MHz): 3.44 p.p.m. (*s*, 8H).

### Crystal data

(C<sub>4</sub>H<sub>12</sub>N<sub>2</sub>)[Cr<sub>2</sub>O<sub>7</sub>]  
*M<sub>r</sub>* = 304.16  
 Monoclinic, *P*2<sub>1</sub>/*c*  
*a* = 11.654 (2) Å  
*b* = 7.8702 (6) Å  
*c* = 10.855 (2) Å  
 $\beta$  = 93.42 (1)°  
*V* = 993.8 (3) Å<sup>3</sup>  
*Z* = 4

*D<sub>x</sub>* = 2.033 Mg m<sup>-3</sup>  
 Mo *K*α radiation  
 Cell parameters from 86 reflections  
 $\theta$  = 26–38°  
 $\mu$  = 2.21 mm<sup>-1</sup>  
*T* = 293 (2) K  
 Block, orange  
 0.13 × 0.11 × 0.07 mm

### Data collection

Stoe AED-II four-circle diffractometer  
 $\omega$ – $\theta$  scans  
 Absorption correction: numerical (*X-SHAPE*; Stoe & Cie, 1998)  
 $T_{\min}$  = 0.755,  $T_{\max}$  = 0.860  
 3793 measured reflections  
 2901 independent reflections  
 2459 reflections with  $I > 2\sigma(I)$

$R_{\text{int}}$  = 0.033  
 $\theta_{\text{max}}$  = 30.0°  
 $h$  = -16 → 16  
 $k$  = -2 → 11  
 $l$  = -15 → 0  
 4 standard reflections  
 frequency: 120 min  
 intensity decay: none

Refinement

Refinement on $F^2$	$w = 1/[\sigma^2(F_o^2) + (0.044P)^2 + 0.3618P]$
$R[F^2 > 2\sigma(F^2)] = 0.026$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.078$	$(\Delta/\sigma)_{\max} = 0.001$
$S = 1.07$	$\Delta\rho_{\max} = 0.51 \text{ e } \text{\AA}^{-3}$
2901 reflections	$\Delta\rho_{\min} = -0.62 \text{ e } \text{\AA}^{-3}$
137 parameters	Extinction correction: <i>SHELXL97</i>
H-atom parameters constrained	Extinction coefficient: 0.0126 (13)

**Table 1**  
Selected geometric parameters ( $\text{\AA}$ ,  $^\circ$ ).

Cr1—O1	1.6068 (14)	N1—C2	1.487 (2)
Cr1—O3	1.6101 (14)	N1—C1	1.498 (2)
Cr1—O2	1.6334 (14)	C1—C2 <sup>i</sup>	1.512 (2)
Cr1—O4	1.7890 (13)	C2—C1 <sup>i</sup>	1.512 (2)
Cr2—O5	1.5939 (16)	N2—C3	1.484 (3)
Cr2—O6	1.6189 (16)	N2—C4	1.490 (3)
Cr2—O7	1.6242 (14)	C3—C4 <sup>ii</sup>	1.504 (3)
Cr2—O4	1.7830 (13)	C4—C3 <sup>ii</sup>	1.504 (3)
O1—Cr1—O3	110.06 (8)	O6—Cr2—O4	108.91 (7)
O1—Cr1—O2	108.25 (8)	O7—Cr2—O4	107.67 (7)
O3—Cr1—O2	111.67 (8)	Cr2—O4—Cr1	117.79 (7)
O1—Cr1—O4	106.69 (7)	C2—N1—C1	111.69 (13)
O3—Cr1—O4	109.60 (7)	N1—C1—C2 <sup>i</sup>	110.06 (14)
O2—Cr1—O4	110.44 (7)	N1—C2—C1 <sup>i</sup>	110.47 (14)
O5—Cr2—O6	111.42 (11)	C3—N2—C4	111.69 (15)
O5—Cr2—O7	110.45 (8)	N2—C3—C4 <sup>ii</sup>	110.97 (15)
O6—Cr2—O7	109.88 (8)	N2—C4—C3 <sup>ii</sup>	109.77 (15)
O5—Cr2—O4	108.41 (9)		

Symmetry codes: (i)  $2 - x, 2 - y, 1 - z$ ; (ii)  $1 - x, 1 - y, -z$ .

**Table 2**  
Hydrogen-bonding geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N1—H1N1 $\cdots$ O2	0.90	2.08	2.878 (2)	148
N1—H1N1 $\cdots$ O5 <sup>iii</sup>	0.90	2.36	2.925 (2)	121
N1—H2N1 $\cdots$ O2 <sup>iv</sup>	0.90	2.04	2.893 (2)	159
N1—H2N1 $\cdots$ O1 <sup>iv</sup>	0.90	2.56	3.227 (2)	132
N2—H1N2 $\cdots$ O7 <sup>v</sup>	0.90	1.93	2.828 (2)	174
N2—H2N2 $\cdots$ O6	0.90	1.91	2.779 (2)	162

Symmetry codes: (iii)  $x, \frac{3}{2} - y, \frac{1}{2} + z$ ; (iv)  $2 - x, \frac{1}{2} + y, \frac{3}{2} - z$ ; (v)  $x, \frac{3}{2} - y, z - \frac{1}{2}$ .

The C—H and N—H H atoms were positioned with idealized geometry (C—H = 0.97  $\text{\AA}$  and N—H = 0.90  $\text{\AA}$ ) and refined with fixed isotropic displacement parameters according to the riding model [ $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{methylene C,N})$ ].

Data collection: *DIF4* (Stoe & Cie, 1992); cell refinement: *DIF4*; data reduction: *REDU4* (Stoe & Cie, 1992); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *XP* in *SHELXTL* (Bruker, 1998); software used to prepare material for publication: *CIFTAB* in *SHELXTL*.

This work was supported by the state of Schleswig-Holstein and the Deutsche Forschungsgemeinschaft. BRS thanks the Department of Science and Technology (DST), New Delhi, and the Deutscher Akademischer Austauschdienst (DAAD) Bonn for a visiting fellowship under the DST-DAAD (PPP-2004) project.

References

Bruker (1998). *SHELXTL*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 House, H. O. (1977). *Modern Synthetic Reactions*, 2nd ed., pp. 257–266. California: W. A. Benjamin Inc.  
 Movassagh, B., Lakouraj, M. M. & Ghodrati, K. (2002). *Indian J. Chem. Sect. B*, **41**, 1293–1295.  
 Norquist, A. J., Heier, K. R., Halasyamani, P. S., Stern, C. L. & Poeppelmeier, K. R. (2001). *Inorg. Chem.* **40**, 2015–2019.  
 Panagiotopoulos, N. Ch. & Brown, I. D. (1972). *Acta Cryst.* **B28**, 1352–1357.  
 Santaniello, E., Milani, F. & Casati, R. (1983). *Synthesis*, pp. 749–750.  
 Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.  
 Srinivasan, B. R., Dhuri, S. N., Näther, C. & Bensch, W. (2003a). *Acta Cryst.* **C59**, m124–m127.  
 Srinivasan, B. R., Dhuri, S. N., Näther, C. & Bensch, W. (2003b). *Acta Cryst.* **E59**, m639–m641.  
 Srinivasan, B. R., Dhuri, S. N., Näther, C. & Bensch, W. (2003c). *Indian J. Chem. Sect. A*, **42**, 2735–2741.  
 Srinivasan, B. R., Dhuri, S. N., Poisot, M., Näther, C. & Bensch, W. (2004). *Z. Naturforsch.* In the press.  
 Stoe & Cie (1992). *DIF4* (Version 7.09X/DOS) and *REDU4* (Version 7.03). Stoe & Cie, Darmstadt, Germany.  
 Stoe & Cie (1998). *X-SHAPE*. Version 1.03. Stoe & Cie, Darmstadt, Germany.  
 Tran Qui, D. & Palacios, E. (1990). *Acta Cryst.* **C46**, 1212–1215.