

***Camarosporium*-Like Species are Polyphyletic in *Pleosporales*;
Introducing *Paracamarosporium* and *Pseudocamarosporium*
gen. nov. in *Montagnulaceae***

Author(s): Nalin N. Wijayawardene, Kevin D. Hyde, D. Jayarama Bhat, Erio Camporesi, René K. Schumacher, K.W. Thilini Chethana, Saowanee Wikee, Ali H. Bahkali & Yong Wang

Source: *Cryptogamie, Mycologie*, 35(2):177-198. 2014.

Published By: Association des Amis des Cryptogames

DOI: <http://dx.doi.org/10.7872/crym.v35.iss2.2014.177>

URL: <http://www.bioone.org/doi/full/10.7872/crym.v35.iss2.2014.177>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Camarosporium*-like species are polyphyletic in *Pleosporales*; introducing *Paracamarosporium* and *Pseudocamarosporium* gen. nov. in *Montagnulaceae

Nalin N. WIJAYAWARDENE^{a,b}, Kevin D. HYDE^{b,c,d,e}, D. Jayarama BHAT^f,
Erio CAMPORESI^g, René K. SCHUMACHER^h, K.W. Thilini CHETHANA^b,
Saowanee WIKEE^b, Ali H. BAHKALI^e & Yong WANG^{a*}

^aDepartment of Plant Pathology, Agriculture College,
Guizhou University, 550025, P.R. China

^bInstitute of Excellence in Fungal Research and School of Science,
Mae Fah Luang University, Chiang Rai, 57100, Thailand

^cKey Laboratory for Plant Diversity and Biogeography of East Asia,
Kunming Institute of Botany, Chinese Academy of Sciences, 132 Lanhei Road,
Kunming 650201, China

^dWorld Agroforestry Centre, East Asia Office, Kunming 650201, Yunnan, China

^eCollege of Science, Botany and Microbiology Department,
King Saud University, Riyadh 1145, Saudi Arabia

^fFormerly, Department of Botany, Goa University, Goa 403 206, India

^gA.M.B. Gruppo Micologico Forlivese “Antonio Cicognani”,
Via Roma 18, Forlì, Italy; A.M.B. Circolo Micologico “Giovanni Carini”, C.P. 314,
Brescia, Italy; Società per gli Studi Naturalistici della Romagna, C.P. 144,
Bagnacavallo (RA), Italy

^hHölderlinsraße 25, 15517 Fürstenwalde/Spree, Germany

Abstract – *Camarosporium* is a large coelomycetous genus which was formerly recognised as an asexual state in *Botryosphaerales* and *Cucurbitariaceae*. In the present study, we collected several *Camarosporium*-like taxa in Europe (Germany and Italy) and carried out morpho-molecular analyses. Molecular analyses (maximum likelihood, maximum parsimony and MrBayes) of combined LSU and SSU gene datasets show that the *Camarosporium*-like taxa are polyphyletic in *Pleosporales*. *Camarosporium quaternatum*, the type species of *Camarosporium* clusters in the suborder *Pleosporinae* with five other *Camarosporium* species. This clade is supported by high bootstrap and PP values and is distinct from other well-established families in *Pleosporinae*. Other *Camarosporium*-like taxa grouped in *Montagnulaceae* (*Massarineae*) as two phylogenetically distinct clades and are introduced as two new genera, viz. *Paracamarosporium* and *Pseudocamarosporium*. *Paracamarosporium* is morphologically distinct as it has paraphyses and microconidia, while *Pseudocamarosporium* lacks both of these characters. Since *Camarosporium* comprises a large number of species epithets, re-collection and morpho-molecular studies of other *Camarosporium*-like taxa is essential.

Coelomycetes / Massarineae / Multi-gene analyses / Phylogeny / Pleosporinae

* Corresponding author: Yong Wang (email: yongwangbis@aliyun.com)

INTRODUCTION

Camarosporium Schulzer was established by Schulzer (1870) based on *C. quaternatum* (Hazsl.) Schulz. and presently comprises more than 500 species epithets (Index Fungorum, 2014). Members of *Camarosporium* spp. have a worldwide distribution and commonly inhabit branches and leaves as saprobes of a wide range of hosts (Sutton, 1980; Farr & Rossman, 2014). Taylor *et al.* (2001) reported on *Camarosporium* species associated with leaf spots of *Protea cynaroides* but suggested they were not economically important phytopathogens. However, *Camarosporium* has received attention in taxonomic studies as the genus was considered heterogeneous in morphology (Sutton, 1980). Taxonomic notes of *Camarosporium* and other muriform genera (such as *Coryneum*, *Dichomera*), based on morphology, have been undertaken by Sutton & Pollack (1974) and Sutton (1975, 1977, 1980). With the implementation of molecular techniques in fungal taxonomy profound changes have taken place in our understanding of the fungi and are resulting in a natural classification based on morphology as well as molecular data (Chomnunti *et al.*, 2011; Boonmee *et al.*, 2012; Wijayawardene *et al.*, 2014a). Molecular studies on the phylogenetic placement of the genus and species of *Camarosporium* have been carried out by Wijayawardene *et al.*, (2014b) and showed that *Camarosporium sensu stricto* belongs to *Pleosporinae* (*Pleosporales*). Since there are several muriform coelomycetes which are morphologically similar to *Camarosporium* (Sutton, 1980) it is important to resolve generic boundaries based on molecular data analyses and incorporate them into a natural classification system (Wijayawardene *et al.*, 2013c).

We collected several *Camarosporium*-like species from Europe (Germany and Italy) and these were subjected to morpho-molecular study. Mega blast searches of LSU and SSU sequence data showed three species were linked with *Camarosporium quaternatum*, the type species of *Camarosporium* (in *Pleosporinae*, *Pleosporales*), while other strains were related to *Montagnulaceae*, *Massarineae*. In this paper, the placement of *Camarosporium sensu stricto* in *Pleosporinae* is confirmed and the six other *Camarosporium*-like species are placed in the new genera *Paracamarosporium* and *Pseudocamarosporium* in *Montagnulaceae* (*Massarineae*, *Pleosporales*) which are introduced herein, based on molecular and morphological study.

MATERIALS AND METHODS

Collecting, morphological studies and isolation: Fresh decaying plant materials were collected from several localities in Germany and Italy, placed in paper bags and returned to laboratory. Conidiomata on fresh samples were observed under a stereomicroscope and removed using a sterilised needle and placed in a droplet of distilled water on a clean slide. Conidial structures (i.e. conidiophores, conidiogenous cells and conidia) were examined under a compound microscope (Nikon Eclipse 80i compound microscope fitted with a Canon 450D digital camera). Single conidial isolation was carried out as described in Chomnunti *et al.* (2011). Germinating conidia were transferred aseptically to potato dextrose agar (PDA) plates and grown at 18°C. Colony colour and other characters were

assessed after 1 week. The specimens are deposited in the herbarium of Mae Fah Luang University (MFLU), Chiang Rai, Thailand, New Zealand Fungal & Plant Disease Collection, Landcare Research, New Zealand (PDD) and Guizhou University Fungi Collection (GUFC). Living cultures are deposited at the Culture Collection at Mae Fah Luang University (MFLUCC), Landcare Research, New Zealand (ICPM) and Department of Plant Pathology, Agriculture College, Guizhou University, China (GUCC).

DNA extraction, PCR amplification and sequencing: Colonies generated from single spores were grown on PDA for 7-14 days at 18°C. Fresh fungal mycelia were used to extract genomic DNA by using a BIOMIGA Fungus Genomic DNA Extraction Kit (GD2416) (Wijayawardene *et al.*, 2013a, b). The amplification of rDNA regions of the internal transcribed spacers (ITS), small subunit rDNA (SSU) and large subunit (LSU) genes was carried out by using ITS5 and ITS4, NS1 and NS4 and LROR and LR5 (White *et al.*, 1990; Vilgalys & Hester, 1990) primers. Optimum conditions for amplification of ITS and LSU regions are as described in Alves *et al.* (2004, 2005) and for SSU region as described in Phillips *et al.* (2008). Amplified PCR fragments were checked on 1% agarose electrophoresis gels stained with ethidium bromide. Purified PCR products (by minicolumns, purification resin and buffer according to the manufacturer's protocols Amersham product code: 27-9602-01) were sent to SinoGenoMax Co., Beijing, China for DNA sequencing. The nucleotide sequence data obtained are deposited in GenBank (Table 1).

Table 1. Strains used in this study. Type strains are in bold and newly generated sequences are in bold and marked with an asterisk

Taxon	Culture collection number	GenBank Accession number		
		LSU	SSU	ITS
<i>Alternaria alternata</i>	CBS 916.96		KC584507	
<i>Ampelomyces quisqualis</i>	CBS 129.79	EU754128	EU754029	
<i>Bambusicola bambusae</i>	MFLUCC 11-0614	JX442035	JX442039	JX442031
<i>Bambusicola massarinia</i>	MFLUCC 11-0389	JX442037	JX442041	JX442033
<i>Bambusicola splendida</i>	MFLUCC 11-0439	JX442038	JX442042	JX442034
<i>Bimuria novae-zelandiae</i>	CBS 107.79	AY016356	AY016338	
<i>Camarosporium aloes</i>	CPC 21572	KF777198		
<i>Camarosporium clematidis</i>	MFLUCC 13-0336	KJ724249		
<i>Camarosporium elongata</i>	AFTOL-ID 1568	DQ678061	DQ678009	
<i>Camarosporium elongata</i>	MFLUCC 14-0260	KJ724249		
<i>Camarosporium quaternatum</i>	CBS 483.95	GU301806	GU296141	
<i>Camarosporium robinium</i>	MFLUCC 13-0527			
<i>Camarosporium</i> sp.	CPC 12441	DQ377885		
<i>Camarosporium spartii</i>	MFLUCC 13-0548			
<i>Cochliobolus heterostrophus</i>	AFTOL-ID 54	AY544645	AY544727	
<i>Cochliobolus sativus</i>	AFTOL-ID 271	DQ678045	DQ677995	
<i>Coniothyrium palmarum</i>	CBS 400.71	EU754153	EU754054	
<i>Coniothyrium palmarum</i>	CBS 758.73	EU754154	EU754055	
<i>Cucurbitaria berberidis</i>	CBS 394.84	GQ387605	GQ387544	

Table 1. Strains used in this study. Type strains are in bold and newly generated sequences are in bold and marked with an asterisk (*continued*)

Taxon	Culture collection number	GenBank Accession number		
		LSU	SSU	ITS
<i>Cucurbitaria berberidis</i>	MFLUCC 11-0384	KC506793		
<i>Cucurbitaria berberidis</i>	MFLUCC 11-0385	KC506794		
<i>Decorospora gaudefroy</i>	CBS 332.63	EF177849	AF394542	
<i>Deniquelata barringtoniae</i>	MFLUCC 110422			
<i>Didymella exigua</i>	CBS 183.55	JX681089	GU296147	GU237794
<i>Didymella pisi</i>	CBS 126.54	GU237968	EU754038	
<i>Didymocrea sadasivani</i>	CBS 438.65	DQ384103	DQ384074	
<i>Dothidothia aspera</i>	CPC 12933	EU673276	EU673228	
<i>Dothidothia symphoricarpi</i>	CPC 12929	EU673273	EU673224	
<i>Halomassarina thalassiae</i>	JK 5385B	GU479804		
<i>Halomassarina thalassiae</i>	JK 5262D	GU301816		
<i>Helicascus nypae</i>	BCC 36752	GU479789	GU479755	
<i>Julella avicenniae</i>	BCC 20173	GU371822	GU371830	
<i>Julella avicenniae</i>	BCC 18422	GU371823	GU371831	
<i>Kalmusia brevispora</i>	KT 1466	AB524600	AB524459	
<i>Kalmusia ebuli</i>	CBS 123120	JN644073	JN851818	
<i>Karstenula rhodostoma</i>	CBS 690.94	GU301821	GU296154	
<i>Lentithecium aquaticum</i>	CBS 123099	GU301823	GU296156	
<i>Lentithecium arundinaceum</i>	CBS 619.86	GU301824	GU296157	
<i>Lentithecium fluviatile</i>	CBS 122367	GU301825	GU296158	
<i>Leptosphaeria doliolum</i>	CBS 541.66	JF740284		
<i>Leptosphaeria doliolum</i>	CBS 155.94	JF740282		
<i>Leptosphaeria doliolum</i>	CBS 125979	JF740283		
<i>Leptosphaeria errabunda</i>	CBS 617.75	JF740289		
<i>Leptosphaeria sydowii</i>	CBS 385.80	JF740313		
<i>Leptosphaerulina australis</i>	CBS 317.83	GU301830	GU296160	
<i>Massarina cisti</i>	CBS 266.62	FJ795447	FJ795490	
<i>Massarina eburnea</i>	CBS 473.64	GU301840	GU296170	
<i>Montagnula anthostomoides</i>	CBS 615.86	GU205223	GU205246	
<i>Montagnula anthostomoides</i>	CBS 615.86	GU205223	GU205246	
<i>Montagnula opulenta</i>	CBS 168.34	NG_027581		
<i>Morosphaeria ramunculicola</i>	JK 5304B	GU479794	GU479760	
<i>Neottiosporina paspali</i>	CBS 331.37	EU754172	EU754073	
<i>Ophiosphaerella herpotricha</i>	AFTOL-ID 1595	DQ767656	DQ767650	
<i>Paracamarosporium psoraleae</i>	CPC 21632	KF777199		KF777143
<i>Paraconiothyrium estuarinum</i>	CBS 109850		AY642522	AY642530
<i>Paraconiothyrium fuscomaculans</i>	CBS 116.16	EU754197	EU754098	
<i>Paraconiothyrium minitans</i>	CBS 122786	EU754174	EU754075	
<i>Paraconiothyrium minitans</i>	CBS 122788	EU754173	EU754074	

Table 1. Strains used in this study. Type strains are in bold and newly generated sequences are in bold and marked with an asterisk (*continued*)

Taxon	Culture collection number	GenBank Accession number		
		LSU	SSU	ITS
<i>Paraphaeosphaeria michotii</i>	CBS 652.86	GQ387581	GQ387520	
<i>Paraphaeosphaeria michotii</i>	CBS 652.86	GQ387581	GQ387520	
<i>Paraphaeosphaeria michotii</i>	CBS 591.73	GU456326	GU456305	
<i>Paraphoma fimeti</i>	CBS 170.70	GQ387584	GQ387523	
<i>Peyronellaea zeae-maydis</i>	CBS 588.69	EU754192	EU754093	
<i>Phaeosphaeria nodorum</i>	CBS 110109	EU754175	EU754076	
<i>Phaeosphaeria oryzae</i>	CBS 110110	GQ387591	GQ387530	
<i>Phoma herbarum</i>	CBS 615.75	EU754186	EU754087	
<i>Pleospora calvescens</i>	CBS 246.79	EU754131	EU754032	
<i>Pleospora herbarum</i>	CBS 191.86	GU238160	GU238232	
<i>Pleospora typhicola</i>	CBS 132.69	JF740325	JF740105	
<i>Pseudocamarosporium corni</i>*	MFLUCC 13-0541	KJ813279*	KJ819946*	KJ747048*
<i>Pseudocamarosporium lonicerae</i>*	MFLUCC13-0532	KJ813278*	KJ819947*	KJ747047*
<i>Pseudocamarosporium piceae</i>*	MFLUCC14-0192	KJ803030*	KJ819948*	KJ747046*
<i>Pseudocamarosporium propinquum</i>*	MFLUCC 13-0544	KJ813280*	KJ819949*	KJ747049*
<i>Pseudocamarosporium tilicola</i>*	MFLUCC 13-0550	KJ813281*	KJ819950*	KJ747050*
<i>Pyrenochaeta acicola</i>	CBS 122789	EU754204	EU754204	
<i>Pyrenochaeta nobilis</i>	CBS 407.76	EU754206	EU754107	
<i>Pyrenochaeta quercina</i>	CBS 115095	GQ387619	GQ387558	
<i>Pyrenochaetopsis decipiens</i>	CBS 343.85	GQ387624	GQ387563	
<i>Pyrenochaetopsis leptospora</i>	CBS 101635	GQ387627	GQ387566	
<i>Pyrenophora phaeocomes</i>	AFTOL-ID 283	DQ499596		
<i>Pyrenophora phaeocomes</i>	AFTOL-ID 283	DQ499596	DQ499595	
<i>Pyrenophora tritici-repentis</i>	AFTOL-ID 173	AY544716	AY544672	
<i>Trematosphaeria pertusa</i>	CBS 122368	FJ201990	FJ201991	
<i>Wojnowicia hirta</i>	CBS 160.73	EU754222	EU754123	

Phylogenetic analyses: A blast search was carried out to reveal the closest taxa to our strains. The sequences were downloaded from GenBank and aligned separately using Bioedit (Hall, 2004), ClustalX v. 1.83 (Thompson *et al.*, 1997) and MEGA 5 (Tamura *et al.*, 2011). Further improvements of the data set were carried out in MAFFTv6 (Kato *et al.*, 2002; Kato & Toh, 2008), online sequence alignment editor under the default settings (mafft.cbrc.jp/alignment/server/) and alignments were checked and manual adjustments were made wherever necessary. Then the individual datasets finally combined into one dataset. One hundred thorough maximum likelihood was performed in RAxML 7.2.8 as part of the “RAxML-HPC2 on TG” tool (Stamatakis, 2006) implemented in raxmlGUI v.0.9b2 (Silvestro & Michalak, 2010). Maximum-parsimony (MP) analysis was carried out using PAUP v. 4.0b10 (Swofford, 2002). Posterior probabilities (PP) (Rannala & Yang, 1996; Zhaxybayeva & Gogarten, 2002) were valued by Markov Chain Monte Carlo sampling (BMCMC) in MrBayes v. 3.0b4 (Huelsenbeck & Ronquist, 2001).

RESULTS

Phylogenetic analyses: Combined analyses of LSU and SSU dataset of families in suborders of *Pleosporales* i.e. *Pleosporinae* (*Coniothyriaceae*, *Cucurbitariaceae*, *Didymellaceae*, *Dothidotthiaceae*, *Leptosphaeriaceae*, *Phaeosphaeriaceae* and *Pleosporaceae*) and *Massarineae* (*Bambusicolaceae*, *Lentitheciaceae*, *Massarinaceae*, *Montagnulaceae*, *Morosphaeriaceae*, *Julellaceae* and *Trematosphaeriaceae*) (Zhang *et al.*, 2012; Hyde *et al.*, 2013) were used to carry out phylogenetic analyses (Table 1). A separate data set which comprised LSU, SSU and ITS sequence data was used to show the placement of new genera and new species in *Montagnulaceae*. A heuristic search with random addition of taxa (1000 replicates) and gaps are treated as missing characters. Bootstrap support (BS) values of MP and ML (equal or above 50 %) are shown on the upper branches and PP values (equal or above 0.7) from MCMC analyses are shown under the branches.

Combined gene (LSU and SSU) analyses for Pleosporinae: The combined LSU and SSU data set consists of 43 taxa with the outgroup taxa i.e. *Corynespora cassicola* (CBS 100822) and *C. smithii* (CABI 5649b). The data set consists of 2683 (LSU = 1285 bp and 1390 bp) characters after alignment of which 2642 are included in ML, MP and Bayesian analyses. We have carried out Bayesian analysis for *Pleosporinae* group as some nodes are not supported with higher bootstrap values in MP and ML analyses. Of the included bases, 2350 sites are conserved regions while 292 and 160 sites are variables and parsimony informative respectively.

Camarosporium quaternatum (CBS 483.95), the type species of *Camarosporium* groups with other *Camarosporium* spp. viz. *C. aloes* (CPC 21572 *fide* Crous *et al.*, 2013), *C. clematidis* (MFLUCC 13-0336 *fide* Wijayawardene *et al.*, 2014b), *C. robinicola* (MFLUCC 13-0527 *fide* Wijayawardene *et al.*, 2014b), *C. spartii* (MFLUCC 13-0548 *fide* Wijayawardene *et al.*, 2014b) and *Camarosporium* sp. (CPC 12441 *fide* Crous *et al.*, 2006) in a separate clade. This clade is supported by moderate bootstrap values (75% in ML analysis) and high PP value (1.00). This clade is distinct from *Cucurbitariaceae* and *Leptosphaeriaceae* where *Camarosporium* was previously stated to be the asexual state (Schoch *et al.*, 2009; Doilom *et al.*, 2013). *Cucurbitaria elongata* (AFTOL-ID 1568 *fide* Schoch *et al.*, 2006; MFLUCC 14-0260 *fide* Wijayawardene *et al.*, 2014b) however, also cluster in the *Camarosporium* clade and thus is introduced as a new combination.

Combined gene (LSU and SSU) analyses for Massarineae: The combined LSU and SSU data set consists of 27 taxa with *Pleospora herbarum* (CBS 191.86) as the outgroup taxon. The data set consists of 2819 characters (LSU = 1310 bp and SSU = 1500 bp) after alignment of which 2735 are included in ML and MP analyses. Of the included bases, 2191 sites are conserved regions, while 544 and 261 sites are variables and parsimony informative respectively.

Our new collections of *Camarosporium*-like taxa and *Camarosporium psoraleae* (CPC 21632 *fide* Crous *et al.*, 2013) cluster in *Montagnulaceae* (Fig. 2).

Combined gene (LSU, SSU and ITS) analyses for Montagnulaceae: The combined LSU, SSU and ITS data set used to show the generic placement of *Montagnulaceae*. The data set consists of 22 taxa with *Julella avicenniae* (BCC 18422 and BCC 20173) as the outgroup taxon. The data set consists of 3330 characters (LSU = 1284 bp, SSU = 1501 bp and ITS = 545 bp) after alignment of which 2880 are included in ML and MP analyses. Of the included bases,

2525 sites are conserved regions, while 355 and 175 sites are variables and parsimony informative respectively.

Clade 1 in Figure 3 comprises our five strains of *Camarosporium*-like taxa and this clade is not related with *Camarosporium sensu stricto*. Hence *Pseudocamarosporium* which has high bootstrap values (ML: 99% and MP: 100%) is introduced to accommodate these taxa. *Pseudocamarosporium* comprises with five species viz. *P. corni*, *P. lonicerae*, *P. piceae*, *P. propinquum* (\equiv *Camarosporium propinquum* (Sacc.) Sacc.) and *P. tillicola*.

Camarosporium psoraleae (CPC 21632 *vide* Crous *et al.*, 2013) distinct from *Camarosporium sensu stricto* (*Pleosporinae*) is placed in *Montagnulaceae* (Clade 1, Fig. 3). However, it has a distinct phylogenetic lineage from *Pseudocamarosporium*, hence we introduce *Paracamarosporium* as a new genus.

TAXONOMY

In multi-gene analyses of *Pleosporinae* (Fig. 1), *Cucurbitaria elongata* (AFTOL-ID 1568 *vide* Schoch *et al.*, 2006; MFLUCC 14-0260) cluster in *Camarosporium sensu stricto* clade.

Camarosporium elongata* (Fr.) Grev. Ex. Wijayawardene & K.D. Hyde *comb. nov.

Index Fungorum Number: IF550571

\equiv *Cucurbitaria elongata* (Fr.) Grev., Scott. crypt. fl. (Edinburgh) 4(37-48): pl. 195 (1826) [1825]

= *Cucurbitaria elongata* var. *coronillae* Fuckel, Jb. Nassau. Ver. Naturk. 23-24: 174 (1870) [1869-70]

= *Cucurbitaria elongata* (Fr.) Grev., Scott. crypt. fl. (Edinburgh) 4(37-48): pl. 195 (1826) [1825] var. *elongata*

= *Gibberidea elongata* (Fr.) Kuntze, Revis. gen. pl. (Leipzig) 3(2): 481 (1898)

= *Sphaeria elongata* Fr., Observ. mycol. (Havniae) 1: 175 (1815)

Illustrations: Hyde *et al.*, 2013 (Figure 32, a-i, illustrated as *Cucurbitaria elongata* (Fr.) Grev.)

Notes: Our molecular data analyses (Figs 1 & 2) show *Camarosporium*-like taxa are polyphyletic in *Pleosporales*. *Camarosporium sensu stricto* groups in *Pleosporinae* while other taxa group as two distinct sub-clades in *Massarineae*. One clade comprises five species hence *Pseudocamarosporium* is introduced to accommodate them. The second clade comprises one strain thus *Paracamarosporium* is introduced to accommodate it.

Paracamarosporium* Wijayawardene & K.D. Hyde *gen. nov.

Index Fungorum Number: IF550563

Etymology: Named after its morphological similarity to the genus *Camarosporium*

Saprobic on dead branches and stems of *Psoralea pinnata* (*Fabaceae*).

Sexual state: Not observed. **Asexual state**: *Conidiomata* immersed to erumpent, solitary, globose, with wall of 3-6 layers of brown cells of *textura angularis*, with central ostiole. *Paraphyses* hyaline, hyphae-like, smooth, intermingled among conidiogenous cells, subcylindrical, with bulbous base, tapering to obtuse apex, 1-4-septate, unbranched or branched at base, and anastomosing. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* lining the inner cavity,




Fig. 1. The best scoring RAxML tree from 43 strains based on combined dataset of LSU and SSU sequences. Bootstrap support values for maximum-likelihood (ML) and maximum-parsimony (MP) greater than 50% are given above the nodes. Posterior Probability values (PP) equal to or greater than 0.7 are given below the nodes. The culture collection numbers are given after the species names. The tree is rooted to *Corynespora smithii* (CABI 5649b) and *C. smithiicola* (CBS 100822). Type and extype strains are in bold.

globose to doliiform, hyaline, smooth, phialidic with prominent periclinal thickening and thick channel (at times also with percurrent proliferation). *Conidia* brown, finely roughened, ellipsoid to ovoid, with obtuse ends, 1-3 transversely septate, developing 1-6 oblique to transverse septa, at times becoming constricted at primary septa. *Microconidiogenous* cells intermingled among macroconidiogenous cells, hyaline, smooth, ampulliform to doliiform to irregular, mono- to polyphialidic, proliferating percurrently, or with periclinal thickening. *Microconidia* hyaline, smooth, guttulate, bacilliform to subcylindrical, apex obtuse, base truncate (Description based on Crous *et al.*, 2013).




Fig. 2. The best scoring RAxML tree from 27 strains based on combined LSU and SSU dataset. Bootstrap support values for maximum-likelihood (ML) and maximum-parsimony (MP) values equal to greater than 50% are given above the nodes. The culture collection numbers are given after the species names. The tree is rooted to *Pleospora herbarum* (CBS 191.86). Type and extype strains are in bold and newly introduced taxa are in red.

Notes: *Paracamarosporium* shows similar conidial morphology with *Camarosporium sensu stricto* (Crous *et al.*, 2013), but *Camarosporium sensu stricto* lack paraphyses and microconidia (Sutton, 1980; Wijayawardene *et al.*, 2014b). The molecular data (Fig. 3) shows it is unrelated with *Camarosporium sensu stricto* and clusters in *Montagnulaceae*.

Type species: ***Paracamarosporium psoraleae*** (Crous & M.J. Wingf.) Wijayawardene & K.D. Hyde **comb. nov.**

Index Fungorum Number: IF 550562

≡ *Camarosporium psoraleae* Crous & M.J. Wingf., in Crous *et al.* Persoonia, Mol. Phyl. Evol. Fungi 31: 235 (2013)

Type: South Africa, Western Cape Province, Betty's Bay, Harold Porter National Botanical Garden, on stems of *Psoralea pinnata* (*Fabaceae*), 28. October 2012, M.J. Wingfield (holotype CBS H-21440, culture ex-type CPC 21632 = CBS 136628).

Illustrations: Crous *et al.* 2013 (Fungal Planet 181, illustrated as *Camarosporium psoraleae* Crous & M.J. Wingf.)

Pseudocamarosporium Wijayawardene & K.D. Hyde **gen. nov.**

Index Fungorum Number: IF 550556; *Facesoffungi number:* 00007

Etymology. Named after its morphological similarity to the genus *Camarosporium*




Fig. 3. The best scoring RAxML tree from 22 strains based on LSU, SSU and ITS combined dataset. Bootstrap support values for maximum-likelihood (ML) and maximum-parsimony (MP) values equal to or greater than 50% are given above the nodes. The culture collection numbers are given after the species names. The tree is rooted to *Julella avicenniae* (BCC 18422 and BCC 20173). Type and ex-type strains are in bold and newly introduced taxa are in red (*Pseudocamarosporium*) and blue (*Paracamarosporium*).

Saprobic on dead twigs, branches, cones and stems of various plants. **Sexual state:** Not observed. **Asexual state:** *Conidiomata* pycnidial, black, globose to subglobose, unilocular, immersed, scattered to gregarious, thick-walled, brown to dark brown, with central and papillate ostiole. *Conidiophores* reduced to conidiogenous cell. *Conidiogenous cell* simple, short, hyaline, thin-walled, discrete to integrated, blastic, phialidic with percurrent proliferation. *Conidia* oblong, muriform, with transverse, longitudinal and oblique septa, generally with a truncate base and obtuse apex, varying in shape, brown to dark brown, smooth-walled.

Notes: In conidial morphology, *Pseudocamarosporium* is similar with *Camarosporium sensu stricto* as both genera have muriform conidia and lack pseudoparaphyses. In *Camarosporium*, conidiogenesis is annelidic, while in *Pseudocamarosporium* conidiogenesis is phialidic, but proliferating. Therefore it is quite difficult to distinguish between these two genera based on morphology. *Pseudocamarosporium* is distinct from *Paracamarosporium* in the molecular analyses (Fig. 3) and in morphology. *Paracamarosporium* differs as it has paraphyses and microconidia which are lacking in *Pseudocamarosporium*. We designate *Pseudocamarosporium propinquum* (Sacc.) Sacc. ex. Wijayawardene *et al.* (\equiv *Camarosporium propinquum* (Sacc.) Sacc.) as the type species (Saccardo, 1884).

Pseudocamarosporium corni Wijayawardene, E. Camporesi & K.D. Hyde **sp. nov.**
Fig. 4

Index Fungorum Number: IF 550557; *Facesoffungi number:* 0008

Holotype: MFLU 14-0089

Etymology: Named after the generic name of host, *Cornus*

Saprobic on dead branch of *Cornus sanguinea*. **Sexual state:** Not observed. **Asexual state:** *Conidiomata* pycnidial, 180-270 µm diam., 190-250 µm high, gregarious, dark brown, immersed, unilocular, with centrally located papillate ostiole. *Pycnidial wall* multilayered, outer layer with 1-3 brown walled cells of *textura angularis*, inner wall with 1-4 layers, hyaline. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* enteroblastic, phialidic, percurrently proliferating, smooth, short, hyaline to pale brown. *Conidia* 11-16 × 6-8 µm (\bar{x} = 13.4 × 7.2 µm, n = 20), oblong to ellipsoidal, mostly straight, rarely slight curved, muriform, with 1-3 transverse septa, with 1-2 longitudinal septa, occasionally widest at the middle, brown, smooth-walled.

Culture characteristics: on PDA white from above and very light brown from reverse, with thin mycelium, flat, attaining a diam of 2.5 cm in 7 days at 18°C.

Material examined: Italy, Arezzo Province, Porrena-Poppi, on dead branch of *Cornus sanguinea* L. (*Cornaceae*), 10 March 2013, E. Camporesi, NNW IT 1108 (MFLU 14-0089, **holotype**; isotype PDD 104440), ex-type living culture = MFLUCC 13-0541 = ICMP 20369 = GUCC 0010.




Fig. 4. *Pseudocamarosporium corni* (**holotype**). **A.** Conidiomata on *Cornus sanguinea*. **B, C.** Cross section of pycnidia. **D.** Developing conidium attach to conidiogenous cell. **E.** Conidia. Scale bar: **B, C,** 200 µm; **D, E,** 15 µm.

Notes: Hüseyinov (1968) reported *Camarosporium corni-maris* Hüseyin from *Cornus sanguinea* which has larger conidia ($40\text{--}64 \times 16\text{--}23.5 \mu\text{m}$). Our collection has smaller conidia and molecular data analyses (Figs 2 & 3) show that, it clusters in *Montagnulaceae* and not *Camarosporium sensu stricto* (Fig. 1). Hence we introduce it as a new species in *Pseudocamarosporium*.

Pseudocamarosporium lonicerae Wijayawardene, E. Camporesi & K.D. Hyde
sp. nov. **Fig. 5**

Index Fungorum Number: IF 550558; *Facesoffungi number:* 000018

Etymology: Named after the host, *Lonicera*

Holotype: MFLU 14-0091

Saprobic on stems of *Lonicera* sp. **Sexual state:** Not observed. **Asexual state:** *Conidiomata* pycnidial, 160–180 μm diam., 140–160 μm high, solitary, dark brown, immersed, unilocular, with a papillate ostiole. *Pycnidial wall* multilayered, with 3–4 outer layers of brown-walled cells of *textura angularis*, inner layer with hyaline cells. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* enteroblastic, with percurrent phialidic development, hyaline, smooth, formed on the inner layer of pycnidium wall. *Conidia* $12\text{--}19 \times 5\text{--}7 \mu\text{m}$ ($\bar{x} = 15.3 \times 4.6 \mu\text{m}$, $n = 20$), oblong, mostly straight, rarely slightly curved at the base, muriform, with 3 transverse septa, with 1–2 longitudinal septa, guttulate when young, pale brown to brown at maturity, smooth-walled.

Culture characteristics: on PDA white from above and pale brown from reverse, circular, flat, slow growing, attaining a diam of 2 cm in 7 days at 18°C.

Material examined: Italy, Forlì-Cesena Province, Forlì, Via del Partigiano, on stem of *Lonicera* sp. (*Caprifoliaceae*), 09 March 2013, E. Camporesi, NNW IT 1104 (MFLU 14-0091, **holotype**; isotype PDD 104438), living culture MFLUCC 13-0532 = ICMP 20370 = GUCC 0011.

Notes: Several *Camarosporium* species have been recorded from *Lonicera* sp. viz. *C. caprifolii* Brunaud ($12\text{--}15 \times 5\text{--}6$ fide Brunaud, 1887), *C. lonicerae* S. Ahmad ($22\text{--}26 \times 7.5\text{--}9 \mu\text{m}$ fide Ahmad, 1971), *C. periclymeni* Oudem. ($16\text{--}20 \times 6\text{--}7$ fide Oudemans, 1898), *C. polymorphum* (De Not.) Sacc. (10×8 fide Saccardo, 1884) and *C. xylostei* Sacc. ($18\text{--}20 \times 8$ fide Saccardo, 1884). *Camarosporium caprifolii* has conidia of similar size to our collection, but with only one longitudinal septum (Brunaud, 1887). Molecular data analyses show that our taxon is not congeneric with *Camarosporium sensu stricto* (Figs 1 and 2). Hence we introduce our collection as a new species in *Pseudocamarosporium*.

Pseudocamarosporium piceae Wijayawardene, E. Camporesi & K.D. Hyde
sp. nov. **Fig. 6**

Index Fungorum Number: IF 550559; *Facesoffungi number:* 000019

Etymology: Named after the host genus, *Picea*

Holotype: MFLU 14-0090

Saprobic on cones of *Picea excels.* **Sexual state:** Not observed. **Asexual state:** *Conidiomata* pycnidial, 120–140 μm diam., 110–130 μm high, mostly immersed, unilocular, solitary, scattered, moderately brown, dark brown at ostiolar papilla. *Pycnidial wall* multilayered, with 3–4 outer wall layers of dark brown cells of *textura angularis*, with inner layer with hyaline cells. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* formed from the inner layer of pycnidial wall, enteroblastic, with percurrent phialidic development, smooth, hyaline. *Conidia* $10\text{--}13 \times 6\text{--}7 \mu\text{m}$ ($\bar{x} = 12.2 \times 6.4 \mu\text{m}$, $n = 20$), oblong, muriform,




Fig. 5. *Pseudocamarosporium loniceræ* (holotype). **A**. Conidiomata on *Lonicera* sp. **B, C**. Cross sections of pycnidium. **D, F, G**. Developing conidia attached to conidiogenous cells. **E**. Pycnidium wall. **H-L**. Conidia. **M**. Germinating conidia. Scale bar: **B, C**. 150 μm ; **D, F, G**. 15 μm ; **E**. 75 μm ; **H**. 20 μm , **I-L**. 15 μm ; **M**. 20 μm .

broadly rounded at both ends, with 3 transverse and 1 longitudinal septa, smooth, variable in shape, hyaline when young, dark brown at maturity, smooth-walled.

Cultural characteristics: on PDA white to very light brown from above and reverse, with sparse mycelium, zonate, flat, circular, attaining a diam of 3.5 cm in 7 days at 18°C, reverse become dark green to black in 14 days.




Fig. 6. *Pseudocamarosporium piceae* (**holotype**). **A**. Conidiomata on dead cone of *Picea excels*. **B, C**. Cross sections of pycnidium. **D, E**. Pycnidium wall. **F**. Developing conidia attach to conidiogenous cell. **G-M**. Conidia. Scale bar: **B, C**. 120 μm ; **D, E**. 50 μm ; **F-M**. 10 μm .

Material examined: Italy, Forlì-Cesena Province, San Martino-Predappio, on dead cones of *Picea excels* (*Pinaceae*), 25 March 2012, E. Camporesi, NNW IT 308 (MFLU 14-0090, **holotype**; isotype PDD 104439), living culture MFLUCC 14-0192 = ICMP 20203 = GUCC 0012.

Notes: There are several coelomycetous taxa that have been recorded from *Picea* spp. (Sutton, 1980; Ellis & Ellis, 1985; Farr & Rossman, 2014), but the only *Camarosporium* species recorded is *C. strobilinum* E. Bommer *et al.* However, the conidial dimensions of these two taxa are distinct. *Camarosporium strobilinum* has smaller conidia ($6 \times 3\text{--}3.5 \mu\text{m}$), while *Pseudocamarosporium piceae* has larger conidia ($10\text{--}13 \times 6\text{--}7 \mu\text{m}$).

Pseudocamarosporium propinquum (Sacc.) Wijayawardene, E. Camporesi & K. D. Hyde **comb. nov.**

Fig. 7

≡ *Camarosporium propinquum* (Sacc.) Sacc., Syll. fung. (Abellini) 3: 464 (1884)
 ≡ *Hendersonia propinqua* Sacc., Michelia 1(no. 5): 516 (1879)




Fig. 7. *Pseudocamarosporium propinquum* (epitype). **A.** Conidiomata on *Salix* sp. **B., C.** Cross sections of pycnidia. **D.** Pycnidium wall. **F-H.** Developing conidia attach to conidiogenous cells. **I-Q.** Conidia. **R.** Germinating conidium. Scale bar: **B.** 250 μ m; **C.** 350 μ m; **D.** 50 μ m; **E-H., J-R.** 10 μ m; **I** = 20 μ m.

Index Fungorum Number: IF 550560; *Facesoffungi number*: 000020

Saprobic on dead branch of *Salix sp.* **Sexual state**: Not observed. **Asexual state**: *Conidiomata* pycnidial, 320-400 μm diam., 250-370 μm high, gregarious, black, immersed, unilocular, ostiole papillate, central. *Pycnidial wall* multilayered, with 1-2 outer wall layers of dark brown to black cells of *textura angularis*, inner wall 3-5 layers, hyaline. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* with percurrent phialidic development, smooth, short, hyaline, formed from the inner layer of the pycnidial wall. *Conidia* 11-14 \times 5-7 μm (\bar{x} = 12.2 \times 5.5 μm , n = 20), oblong, with truncate base, apex obtuse, straight, muriform, with 3 transverse septa and 1-2 longitudinal septa, pale brown to dark brown, smooth-walled.

Culture characteristics: on PDA white from above and pale brown from reverse, with thin mycelium, irregular, flat, attaining a diam of 2.5 cm in 7 days at 18°C.

Material examined: FRANCE, Rouen, on branch of *Salix vitellina* L. (*Salicaceae*), Letendre (PAD, holotype); ITALY, Firenze Province, Passo Dell'Eremo – Marradi, on dead branch of *Salix sp.* (*Salicaceae*), 12 May 2013, E. Camporesi, NNW IT 1253 (MFLU 14-0092, **epitype** designated here; isopitype PDD 104441), ex-type living culture MFLUCC 13-0544 = ICMP 20371 = GUCC 0013.



Fig. 8. *Camarosporium propinquum* (holotype).

Notes: Boomer & Rousseau (1884) and Saccardo (1884) described *Camarosporium salicinum* Sacc. *et al.* (conidia $18\text{-}20 \times 8\text{-}10 \mu\text{m}$) and *C. propinquum* (conidia $15\text{-}16 \times 8 \mu\text{m}$) from *Salix vitellina*. Our collection is morphologically similar to *C. propinquum*. Our collection shares similar characters i.e. conidiogenesis and conidial dimensions to the iconotype of *C. propinquum* (Fig. 8). In molecular data analyses, *C. propinquum* clusters in *Montagnulaceae* (Figs 2 & 3). Hence, we transfer this species to *Pseudocamarosporium* as a new combination i.e. *Pseudocamarosporium propinquum* and epitypify *Camarosporium propinquum*.

Pseudocamarosporium tiliicola Wijayawardene, R. K. Schumacher & K. D. Hyde
sp. nov. **Fig. 9**

Index Fungorum Number: IF 550561; *Facesoffungi number:* 000021

Etymology: Named after the host genus *Tilia* from which it was collected

Holotype: MFLU 14-0093

Saprobic in the bark of a dead and corticated branch of *Tilia*. **Sexual state:** Not observed. **Asexual state:** *Conidiomata* pycnidial, 670-700 μm diam., 650-720 μm high, solitary to gregarious, black, semi-immersed to superficial, unilocular, centrally papillate ostiole. *Pycnidial wall* multilayered, with 1-4 outer wall layers of dark brown cells of *textura angularis*, 5-10 cell layers in ostiole




Fig. 9. *Pseudocamarosporium tiliicola* (holotype). **A.** Conidiomata on *Tilia* sp. **B.** Spore mass in cross section. **C, D.** Pycnidium wall. **E, F.** Different stages of developing conidia. **G-K.** Conidia. Scale bar: **B.** 650 μm ; **C.** 300 μm ; **D.** 120 μm ; **E, F.** 15 μm ; **G-K.** 20 μm .

region, inner wall 2-4 layers, hyaline. *Conidiophores* reduced to conidiogenous cells. *Conidiogenous cells* with percurrent phialidic development, smooth, short, hyaline, segregated, formed from the inner layer of the pycnidial wall. *Conidia* 21-26 × 8-12 μm (\bar{x} = 23.8 × 10.2 μm, n = 20), oblong to ellipsoid, straight to curved, rarely rounded, base tapered and seldom distinctly truncate, muriform, with 3-5 transverse septa and occasionally 2-3 longitudinal septa, brown to dark brown, smooth-walled.

Culture characteristics: on PDA white from above and pale brown from reverse, zonate, with thin mycelium, flat, circular, attaining a diam of 3 cm in 7 days at 18°C.

Material examined: Germany, near Berlin, on a branch of *Tilia* sp. (*Malvaceae*), 31 March 2013, Rene K. Schumacher, NNW G3 (G2/9) (MFLU 14-0093, **holotype**; isotype PDD 104442), living culture MFLUCC 13-0550 = ICMP 20372 = GUCC 0014.

Notes: Saccardo (1882) described *Camarosporium tiliae* Sacc. & Penz. from *Tilia europaea* L., however *Pseudocamarosporium tiliicola* has larger conidia (21-26 × 8-12 μm vs 8-10 × 6-7 μm) and is therefore introduced as a new species.

DISCUSSION

New separate clade in Pleosporinae: The *Pleosporales* is one of the most important and significant orders in *Dothideomycetes* (Schoch *et al.*, 2006, 2009; Hyde *et al.*, 2013) and this order has been the subject of morphological (Sivanesan, 1984) and molecular (Zhang *et al.*, 2012) studies. In their molecular based overview of *Dothideomycetes*, Schoch *et al.* (2009) showed that suborder *Pleosporinae* comprises the families *Didymellaceae*, *Leptosphaeriaceae*, *Phaeosphaeriaceae* and *Pleosporaceae*. However, Zhang *et al.* (2012) predicted the possibilities of introducing *Cucurbitariaceae* and *Dothidothiaceae* as families in *Pleosporinae*. Hyde *et al.* (2013) agreed with Zhang *et al.* (2012) and showed *Pleosporinae* comprises *Cucurbitariaceae*, *Didymellaceae*, *Dothidothiaceae*, *Leptosphaeriaceae*, *Phaeosphaeriaceae* and *Pleosporaceae* and *Coniothyriaceae* (Hyde *et al.*, 2013).

Conventionally, the genus was linked to *Botryosphaeriales incertae sedis* (Kirk *et al.*, 2008; Liu *et al.*, 2012; Wijayawardene *et al.*, 2012a; Hyde *et al.*, 2013) and *Cucurbitariaceae* (Saccardo, 1884; Sivanesan, 1984; Zhang *et al.*, 2012; Doilom *et al.*, 2013). However, Schoch *et al.* (2009) showed that *Camarosporium quaternatum*, the type species of *Camarosporium* groups in *Leptosphaeriaceae*. Furthermore, Ramaley and Barr (1995) stated that *Pleoseptum yuccaesedum* A.W. Ramaley & M.E. Barr, the type species of *Pleoseptum* A.W. Ramaley & M.E. Barr (*Phaeosphaeriaceae* *fide* Hyde *et al.*, 2013) had asexual states in *Camarosporium* (*viz.* *C. yuccaesedum* Fairm.). Hence this genus is considered to be polyphyletic as predicted by Sutton (1980).

In our multi-gene analyses of *Pleosporinae* (Fig. 1), *Camarosporium quaternatum* groups with *C. aloes*, *C. clematidis*, *C. robiniicola* and *C. spartii* and forms a distinct clade with high bootstrap values in ML analysis (77%) and in high PP value in Bayesian analysis (1.00). Two strains of *Cucurbitaria elongata* (AFTOL-ID 1568 and MFLUCC 14-0260) also group with these *Camarosporium* strains. However, *Cucurbitaria sensu stricto* (*i.e.* *Cucurbitariaceae* *fide* Doilom *et al.*, 2013; Hyde *et al.*, 2013) form a well-supported clade (bootstrap values are 72%

and 78% in ML and MP analyses respectively; 1.00 in Bayesian analysis) including *Cucurbitaria berberidis* (Pers.) Gray, the type species of *Cucurbitaria* (ex-epitype culture MFLUCC11-0384, MFLUCC 11-0385 *fide* Doilom *et al.* 2013, CBS 394.84 *fide* de Gruyter *et al.*, 2009) and *Pyrenochaeta nobilis* De Not., the type species of *Pyrenochaeta* (ex-type strain CBS 407.76 *fide* de Gruyter *et al.*, 2009). *Pyrenochaeta acicola* (CBS 122789 *fide* de Gruyter *et al.*, 2009), *Pyrenochaeta quercina* (CBS 115095 *fide* de Gruyter *et al.*, 2010) and *Pyrenochaetopsis decipiens* (CBS 343.85 *fide* de Gruyter *et al.*, 2010) also clustered in the same clade.

Camarosporium-like taxa are polyphyletic: Recent molecular phylogenetic studies reveal that several coelomycetous genera are polyphyletic, *viz.* *Coniothyrium* and *Phoma* (Verkley *et al.*, 2004, 2014; de Gruyter *et al.*, 2010, 2012). As predicted by Sutton (1980), our molecular based study confirms that *Camarosporium* is also polyphyletic *i.e.* in *Pleosporinae* (*i.e.* *Camarosporium sensu stricto*) and in *Massarineae* (*i.e.* *Montagnulaceae*) (Figs 2 & 3). However, in this study we have not considered the *Camarosporium*-like taxa in *Botryosphaerales* (Wijayawardene *in prep.*), where *Camarosporium* species have previously been linked (Liu *et al.*, 2012; Hyde *et al.*, 2013). Since the type species of *Camarosporium*, *C. quaternatum* groups in *Pleosporinae*, we consider that particular clade as *Camarosporium sensu stricto*. The other *Camarosporium*-like taxa in *Montagnulaceae* group as two distinct lineages and are hence considered as two different genera *viz.* *Paracamarosporium* and *Pseudocamarosporium*.

Based on our multi-gene analyses, we conclude that delimitation of *Camarosporium*-like taxa based on morphological characters such as conidiomatal structure, conidiogenesis and conidial morphology have not been successful. Hence it is important to carry out molecular based identification along with morphological studies to place species in genera.

It is important to re-collect *Camarosporium*-like taxa as this group comprises more than 500 names in both Index Fungorum (2014) and Robert *et al.* (2005). In our study, we have confirmed that these taxa are scattered across *Pleosporales* with distinct phylogenetic lineages. Therefore, further re-collecting, isolation and molecular based studies (Wijayawardene *et al.*, 2012c) are essential to interpret correct generic concepts for *Camarosporium*-like taxa.

Acknowledgements. Nalin N. Wijayawardene and K.W.T. Chethana are indebted to the Mushroom Research Foundation (MRF), Chiang Rai Province, Thailand for providing Postgraduate Scholarship. K.D. Hyde thanks The Chinese Academy of Sciences, project number 2013T2S0030, for the award of Visiting Professorship for Senior International Scientists at Kunming Institute of Botany. Erio Camporesi grateful Giancarlo Lombardi for his invaluable help in the collecting programme and identifying host plants. Nalin N. Wijayawardene thanks to Dhanushka Udayanga for his precious help to carry out molecular analyses. Nalin N. Wijayawardene grateful to Prof. Vadim A. Mel'nik, Laboratory of the Systematics and Geography of Fungi, Komarov Botanical Institute, Russian Academy of Sciences, Professor Popov Street 2, St. Petersburg, 197376, Russia, Dr. P.M. Kirk, Mycology Section, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3DS, UK and Mr. Ken Hudson, Mycology Publications, CABI, Bakeham Lane, Egham, Surrey, TW20 9TY UK for contributing old literature which were important for this article. K.W.T. Chethana would like to thank Dr. Jiye Yan, Institute of Plant and Environment Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing, China for facilitating molecular studies and providing fellowship during the tenure of which this paper was finalised. Yong Wang would like to thank The International Scientific Cooperated Project of Guizhou Province (No[2013]7004). Appreciation is extended to Mae Fah Luang University grant for studying *Dothideomycetes* (no. 56101020032). We thank to Rossella Marcucci, Herbarium Patavinum (PAD) Erbario dell'Università di Padova, Italy, for contributing herbarium materials.

REFERENCES

- AHMAD S., 1971 — Contributions to the fungi of West Pakistan. XI. *Biologia Lahore* 17(2): 67-85.
- ALVES A., CORREIA A., LUQUE J. & PHILLIPS A.J.L., 2004 — *Botryosphaeria corticola*, sp. nov. on *Quercus* species, with notes and description of *Botryosphaeria stevensii* and its anamorph, *Diplodia mutila*. *Mycologia* 96: 598-613.
- ALVES A., PHILLIPS A.J.L., HENRIQUES I. & CORREIA A., 2005 — Evaluation of amplified ribosomal DNA restriction analysis (ARDRA) as a method for the identification of *Botryosphaeria* species. *FEMS Microbiology Letters* 245: 221-229.
- BOMMER E. & ROUSSEAU M., 1884 — Florule mycologique des environs de Bruxelles. Bulletin de la Société Royale de Botanique de Belgique. 23(1): 13-365.
- BOONMEE S., ZHANG Y., CHOMNUNTI P., CHUKEATIROTE E., TSUI C.K.M., BAHKALI A.H. & HYDE K.D., 2011 — Revision of lignicolous *Tubeufiaceae* based on morphological re-examination and phylogenetic analysis. *Fungal Diversity* 51: 63-102.
- BRUNAUD P., 1887 — Champignons à ajouter à la flore mycologique des environs de Saintes. II. Bulletin de la Société Botanique de France. 34: 428-432.
- CHOMNUNTI P., SCHOCH C.L., AGUIRRE-HUDSON B., KO-KO T.W., HONGSANAN S., JONES E.B.G., KODSUEB R., PHOOKAMSAK R., CHUKEATIROTE E., BAHKALI A.H. & HYDE K.D., 2011 — *Capnodiaceae*. *Fungal Diversity* 51: 103-134.
- CROUS P.W., SLIPPERS B., WINGFIELD M.J., RHEEDER J., MARASAS W.F., PHILLIPS A.J.L., ALVES A., BURGESS T., BARBER P. & GROENEWALD J.Z., 2006 — Phylogenetic lineages in the *Botryosphaeriaceae*. *Studies in Mycology* 55: 235-253.
- CROUS P.W., WINGFIELD M.J., GUARRO J., CHEEWANGKOON R. *et al.*, 2013 — Molecular phylogeny and evolution of fungi – Fungal Planet description sheets: 154-213. *Persoonia* 31: 188-296.
- DE GRUYTER J., AVESKAMP M.M., WOUDEBERG J.H.C., VERKLEY G.J.M., GROENEWALD J.Z. & CROUS P.W., 2009 — Molecular phylogeny of *Phoma* and allied anamorph genera: towards a re-classification of the *Phoma* complex. *Mycological Research* 113: 508-519.
- DE GRUYTER J., WOUDEBERG J.H.C., AVESKAMP M.M., VERKLEY G.J.M., GROENEWALD J.Z. & CROUS P.W., 2012 [2013] — Redisposition of *Phoma*-like anamorphs in *Pleosporales*. *Studies in Mycology* 75: 1-36.
- DE GRUYTER J., WOUDEBERG J.H.C., AVESKAMP M.M., VERKLEY G.J.M., GROENEWALD J.Z. & CROUS P.W., 2010 — Systematic reappraisal of species in *Phoma* section *Paraphoma*, *Pyrenochaeta* and *Pleurophoma*. *Mycologia* 102: 1066-1081.
- DOILOM M., LIU J.K., JAKLITSCH W.M., ARIYAWANSA H., WIJAYAWARDENE N.N., CHUKEATIROTE E., ZHANG M., MCKENZIE E.H.C., GEML J., VOGLMAYR H. & HYDE K.D., 2013 — An outline of the family *Cucurbitariaceae*. *Sydowia* 65(1): 167-192.
- ELLIS M.B. & ELLIS J.P., 1985 — Microfungi on land plants: an identification handbook. Richmond Pub. (2nd edition 1997).
- FARR D.F. & ROSSMAN A.Y., 2014 — Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA. Retrieved April 7, 2014, from <http://nt.ars-grin.gov/fungal-databases/>
- HALASZ G., 2001 — Fungi associated with the decline of black locust (*Robinia pseudoacacia* L.) plantations in Hungary. *Journal of Forest Science* 47(2): 88-92.
- HALL T., 2004 — Bioedit version 6.0.7.; Available from: <http://www.mbio.-ncsu.edu/bioedit/bioedit.html>
- HAWKSWORTH D.L., 2012 — Managing and coping with names of pleomorphic fungi in a period of transition. *Mycosphere* 3(2): 143-155.
- HUELSENBECK J.P. & RONQUIST F., 2001 — MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17(8): 754-755.
- HÜSEYINOV E.S., 1968 — [New species of fungi from Azerbaijan]. *Mikologiya i Fitopatologiya* 2(6): 512-514.
- HYDE K.D., JONES E.B.G., LIU J.K., ARIYAWANSA H., BOEHM E. *et al.*, 2013 — Families of *Dothideomycetes*. *Fungal Diversity* 63: 1-313.
- INDEX FUNGORUM 2014 — <http://www.indexfungorum.org/names/NamesRecord.asp?RecordID=489566> (Accession date 02.02.2014).
- KATOH K. & TOH H., 2008 — Recent developments in the MAFFT multiple sequence alignment program. *Briefings in Bioinformatics* 9: 276-285.
- KATOH K., MISAWA K., KUMA K. & MIYATA T., 2002 — MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* 30: 3059-3066.

- KIRK P.M., CANNON P.F., MINTER D.W. & STALPERS J.A., 2008 — Dictionary of the Fungi, 10th edn. CABI, Wallingford.
- KOHLI D.K. & BACHHAWAT A.K., 2003 — CLOURE: Clustal output Reformatter, a program for reformatting ClustalX/ClustalW outputs for SNP analysis and molecular systematics. *Nucleic Acids Research* 31: 3501-3502.
- LIU J.K., PHOOKAMSAK R., DOILOM M., WIKKEE S., LI Y., ARIYAWANSHA H., BHOONMEE S., CHOMNUNTI P., DAI D.Q., BHAT J.D., ROMERO A.I., ZHUANG W.I., MONKAI J., JONES E.B.G., CHUKEATIROTE E., KO KO T.W., ZHAO Y.C., WANG Y. & HYDE K.D., 2012 — Towards a natural classification of *Botryosphaeriales*. *Fungal Diversity* 57: 149-210.
- NAG RAJ T.R., 1993 — Coelomycetous Anamorphs with Appendage-bearing Conidia. Mycologue Publications, Waterloo, Canada.
- OUDEMANS C.A.J.A., 1898 — Contributions á la flore mycologique des Pays-Bas. XVI. *Nederlandsch Kruidkundig Archief* 1: 430-535.
- PAGE R.D.M., 1996 — Tree View: an application to display phylogenetic trees on personal computers. *Computer Applications in the Biosciences* 12: 357-358.
- PHILLIPS A.J.L., ALVES A., PENNYCOOK S.R., JOHNSTON .PR., RAMALEY A., *et al.*, 2008 — Resolving the phylogenetic and taxonomic status of dark-spored teleomorph genera in the *Botryosphaeriaceae*. *Persoonia* 21: 29-55.
- RAMALEY A.W. & BARR M.E., 1995 — New dictyosporous species from leaves of Agavaceae. *Mycotaxon*. 54: 75-90.
- RANNALA B. & YANG Z., 1996 — Probability distribution of molecular evolutionary trees: a new method of phylogenetic inference. *Journal of Molecular Evolution* 43: 304-311.
- ROBERT V., STEGHEUIS G. & STALPERS J., 2005 — The MycoBank engine and related databases. <http://www.mycobank.org> (Accession date: 02.02.2014).
- SACCARDO P.A., 1882 — Fungi Gallici lecti a Cl. viris P. Brunaud, C.C. Gillet, Abb. Letendre, A. Malbranche, J. Therry & Dom. Libert. Series IV. *Michelia*. 2(8): 583-648.
- SACCARDO P.A., 1884 — Sylloge Fungorum III: Patavii, pp 1-815.
- SCHOCH C.L., CROUS P.W., GROENEWALD J.Z., BOEHM E.W.A., BURGESS T.I., DE GRUYTER J. *et al.* 2009 — A class-wide phylogenetic assessment of Dothideomycetes. *Studies of Mycology* 64(1): 1-15.
- SCHOCH C.L., SHOEMAKER R.A., SEIFERT K.A., HAMBLETON S., SPATAFORA J.W. & CROUS P.W. 2006 — A multigene phylogeny of the Dothideomycetes using four nuclear loci. *Mycologia* 98: 1041-1052.
- SCHULZER S., 1870 — Mykologische Beiträge. Verhandlungen der Zoologisch-Botanischen Gesellschaft Wien. 20: 635-658.
- SILVESTRO D. & MICHALAK I., 2010 — raxmlGUI: a graphical front-end for RAxML <http://sourceforge.net/projects/raxmlgui/>. Accessed August 2010.
- SIVANESAN A., 1984 — The bitunicate ascomycetes and their anamorphs. J. Cramer, Vaduz.
- STAMATAKIS A., 2006 — RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22: 2688-2690.
- SUTTON B.C. & POLLACK F.G., 1974 — Microfungi on *Cercocarpus*. *Mycopathologia et mycologia applicatation* 52: 331-351.
- SUTTON B.C., 1975 — Coelomycetes V. *Coryneum*. *Mycological Papers* 138: 1-224.
- SUTTON B.C., 1977 — Coelomycetes VI. Nomenclature of generic names proposed for Coelomycetes. *Mycological Papers* 141: 1-253.
- SUTTON B.C., 1980 — The Coelomycetes. Fungi imperfecti with Pycnidia, Acervuli and Stromata. Commonwealth Mycological Institute, Kew, UK.
- SWOFFORD D.L., 2002 — PAUP: phylogenetic analysis using parsimony, version 4.0 b10. Sinauer Associates, Sunderland MA.
- TAMURA K., PETERSON D., PETERSON N., STECHER G., NEI M. & KUMAR S., 2011 — MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Molecular Biology and Evolution* 28: 2731-2739.
- TAYLOR J.E., CROUS P.W. & SWART L., 2001 — Foliicolous and caulicolous fungi associated with *Proteaceae* cultivated in California. *Mycotaxon* 78: 75-103.
- THOMPSON J.D., GIBSON T.J., PLEWNIAK F., JEANMOUGIN F. & HIGGINS D.G., 1997 — The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 25(24): 4876-4882.
- TRAIL J.W.H., 1888 — Revision of Scotch *Sphaeropsidae* and *Melanconieae*. *Scottish Naturalist* 9: 219-236.
- VERKLEY G.J.M., DUKIK K., RENFURM R., GÖKER M. & STIELOW J.B., 2014 — Novel genera and species of *Coniothyrium*-like fungi in *Montagnulaceae* (Ascomycota). *Persoonia* 32: 25-51.

- VERKLEY G.J.M., SILVA M. D.A., WICKLOW D.T. & CROUS P.W., 2004 — *Paraconiothyrium*, a new genus to accommodate the mycoparasite *Coniothyrium minitans*, anamorphs of *Paraphaeosphaeria*, and four new species. *Studies in Mycology* 50: 323-335.
- VILGALYS R. & HESTER M., 1990 — Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172(8): 4238-4246.
- WHITE T.J., BRUNS T., LEE S. & TAYLOR J., 1990 — Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: PCR Protocols: a guide to methods and applications (Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds). Academic Press, San Diego, California, USA: 315-322.
- WIJAYAWARDENE D.N.N., MCKENZIE E.H.C. & HYDE K.D., 2012a — Towards incorporating anamorphic fungi in a natural classification – checklist and notes for 2011 – *Mycosphere* 3(2): 157-228.
- WIJAYAWARDENE D.N.N., MCKENZIE E.H.C., CHUKEATIROTE E., WANG Y. & HYDE K.D., 2012b — Coelomycetes. *Cryptogamie Mycologie* 33: 215-244.
- WIJAYAWARDENE D.N.N., SONG Y., BHAT D.J., MCKENZIE E.H.C., CHUKEATIROTE E., WANG Y. & HYDE K.D. 2013a — *Wojnowicia viburni* sp. nov. from China and its phylogenetic placement. *Sydowia* 65(1): 181-190.
- WIJAYAWARDENE D.N.N., UDAYANGA D., MCKENZIE E.H.C., WANG Y. & HYDE K.D., 2012c — The future of coelomycete studies. *Cryptogamie Mycologie* 33: 381-391.
- WIJAYAWARDENE N.N., BHAT D.J., HYDE K.D., CAMPORESI E., WIKEE S., CHETHANA K.W.T., TANGTHIRASUNUN N. & WANG Y., 2014b — *Camarosporium sensu stricto* in *Pleosporinae*, *Pleosporales* with two new species. *Phytotaxa* (Submitted).
- WIJAYAWARDENE N.N., CAMPORESI E., BHAT D.J., SONG Y., CHETHANA K.W.T., CHUKEATIROTE E., WANG Y. & HYDE K.D., 2014a — *Macrodidipodopsis* in *Lophiostomataceae*, *Pleosporales*. *Phytotaxa* (in press).
- WIJAYAWARDENE N.N., CAMPORESI E., SONG Y., DAI D.Q., BHAT D.J., MCKENZIE E.H.C., CHUKEATIROTE E., MEL'NIK V.A., WANG Y. & HYDE K.D., 2013b — Multi-gene analyses reveal taxonomic placement of *Scolicosporium minkeviciusii* in *Phaeosphaeriaceae* (*Pleosporales*). *Cryptogamie Mycologie* 34(4): 357-366.
- WIJAYAWARDENE N.N., CROUS P.W., HYDE K.D. & WANG Y., 2013c — The importance of incorporating asexual fungi in to the natural classification system. In: Proceeding of the Asian Mycological Congress, Beijing, China, 2013.
- ZHANG Y., CROUS P.W., SCHOCH C.L. & HYDE K.D., 2012 — *Pleosporales*. *Fungal Diversity* 53: 1-221.
- ZHAXYBAYEVA O. & GOGARTEN J.P., 2002 — Bootstrap, Bayesian probability and maximum likelihood mapping: exploring new tools for comparative genome analyses. *BMC Genomics* 3: 4.