

SF Journal of Environmental and Earth Science

Estuarine Mudflat and Mangrove Sedimentary Environments along Central West Coast of India

Nayak GN* and Noronha e D'Mello CA

Department of Marine Sciences, Goa University, Goa India

Abstract

Estuaries are complex and dynamic aquatic environments, wherein fresh water mixes with rhythmically intruding seawater. Mudflats and mangroves are the important sedimentary sub-environments within estuaries which facilitate deposition of fine sediments, organic matter and metals from overlying water. The cohesive sediments present in these environments protect the coast from erosion and the sediment column can be used to understand past depositional environments, sea level changes, changes in land use-land cover in the catchment area of the rivers and bioavailability of metals.

Keywords: Estuary; Mudflats; Mangroves; Environments

Introduction

The coastal zone is characterized by a variety of landforms out of which estuaries have received considerable attention due to large land-sea interaction mechanisms [1,2]. The word "Estuary" is derived from the Latin "aestus", meaning the tide [3] and Cameron and Pritchard [4] and Pritchard [5] defined an estuary as a semi-enclosed and coastal body of water, with free communication to the ocean, and within which ocean water is diluted by freshwater derived from land. Estuaries are complex and dynamic aquatic environments, where fresh water from river mixes with rhythmically intruding seawater of different physico-chemical composition. Estuaries are major nutrient suppliers to coastal oceans, breeding and nursery grounds for marine organisms and a potential fishery habitat. They act as transportation routes and also recreational places for humans [6,7].

Although all estuaries are analogous, in that they are semi-enclosed bodies of brackish water, a multiple criteria are used to classify them. According to their geological characteristics or geomorphology, estuaries are classified as coastal-plain estuaries, bar-built estuaries or lagoons, fjord-type estuaries, and tectonically caused estuaries [8,9]. Further, based on stratification and circulation, estuaries have also been classified as salt wedge estuaries, vertically homogeneous estuaries and partially mixed estuaries. Further, on the basis of water balance, estuaries are classified into three types: positive, inverse and low-in flow estuaries. Based on tidal range, Hayes [10] defined three types of estuaries: microtidal, mesotidal and macrotidal estuaries.

An estuary is divided into three zones (Figure 1) namely, marine or lower estuary, which has free connections with open sea; middle estuary subjected to strong salt and freshwater mixing and fluvial or an upper estuary, characterized by freshwater but subjected to tidal action. However, the boundaries or the transition zones between these sectors shift according to constantly changing tides and river discharge.

Freshwater inflow plays a key role in carrying continental material from the watershed to the estuary and in balancing effects of tidal inputs of saltwater. Physical, chemical, and biological interactions between terrestrial and coastal systems profoundly affect the transport and fate of material in to the estuary [12]. Material is carried in, from the land via rivers and from the sea by the tides (Figure 2).

Tidal currents provide the steady supply of energy that causes sediment movement into and out of estuaries. Waves and tides carry fine sediments from the mouth of an estuary leaving behind coarser ones. Fresh water from the land drainage adds varying size sediment material during monsoon months. In addition, suspended and particulate material is removed on changes of physicochemical properties on interaction of fresh water and seawater. The distribution of sediments within estuaries is often classified on the basis of grain size which in turn helps in understanding the hydrodynamic conditions prevailing in the area. Estuarine sediments retain large quantity of finer sediments and

OPEN ACCESS

*Correspondence:

Nayak GN, Department of Marine Sciences, Goa University, Goa 403 206, India.

E-mail: gnnayak@unigoa.ac.in

Received Date: 18 Mar 2018

Accepted Date: 23 Apr 2018

Published Date: 30 Apr 2018

Citation: Nayak GN, Noronha e D'Mello CA. Estuarine Mudflat and Mangrove Sedimentary Environments along Central West Coast of India. *SF J Environ Earth Sci.* 2018; 1(1): 1013.

Copyright © 2018 Nayak GN. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1: Estuary showing three divisions—Lower, middle and upper estuary: The boundaries are the transition zones that shift according to season, weather and tides [11].

organic matter and act as sink for a wide range of metals which show high affinity for fine grained sediments. These metals get adsorbed onto the suspended particulate matter and are transported through the water column which finally gets incorporated into the sediments. In this paper the work carried out along central west coast of India on estuarine mudflats and mangroves with an aim to understand past depositional environments, effect of land use-land cover in the catchment area of the rivers and bioavailability of metals and possible bioaccumulation was attempted.

Study area

The west coast of India extends from border with Pakistan in Gujarat to Kanyakumari in Tamil Nadu. It is surrounded by the Arabian Sea. Maharashtra, Goa and Karnataka together comprise the central portion of the west coast of India. This region is highly diverse with respect to coastal environmental features such as estuaries, creeks, lagoons, bays etc. A large number of fast flowing and mostly perennial rivers originate in the Western Ghats and drain into the Arabian Sea. The characteristic of the rivers is the formation of estuaries rather than deltas. Estuaries receive material from multiple sources such as natural weathering of rocks, domestic and industrial sewage outfalls, effluents from mines, and agricultural runoff. The growing population and associated rapid industrialization in and around central west coast of India has increased anthropogenic inputs. Therefore, estuaries are under increasing stress from the human activities.

The catchment area of the rivers in Maharashtra consists of Deccan trap basalts [14] having tholeiitic composition [15]. The catchment area of estuaries in Goa comprises of Western Dharwar Craton (WDC) containing meta-volcanic and meta-sedimentary rocks [16]. The catchment area of estuaries in Karnataka consists of Dharwar system and peninsular gneiss containing metamorphic and crystalline rocks which are made up of granites and granitic-gneisses. The coastline receives the southwest monsoon (June to September) with rainfall decreasing from South (3900mm) to North (2000mm), while the tidal range is in the micro-tidal scale in Karnataka, meso-tidal in Goa and meso to macro tidal in Maharashtra.

Materials and Methods

Sampling is the process or technique of selecting a representative part of a field site in a particular environment for the purpose of determining parameters or characteristics of the entire environment. The sediment cores were collected from mudflat and mangrove environments from almost all the estuaries along central west coast of India covering the coast of Maharashtra, Goa and Karnataka states representing lower, lower middle and upper middle regions estuaries. Precautions were taken to avoid contamination of sediment samples

during collection and handling and a systematic procedure for the analysis of the sediments was followed. The sampling locations were positioned using a hand held Global positioning system (GPS). The sediment cores were collected during the low tide when the intertidal region was well exposed using a pre-cleaned handheld PVC corer. The sediment cores were then sub-sampled at 2cm intervals with the help of plastic knife and transferred into clean and labeled plastic bags with care taken to avoid metal contamination. The packed sub samples were then store in an ice box and transported to the laboratory.

The parameters studied for each sub-sample are pH; sand, silt, clay percentage [17]; clay minerals [18] and were identified and quantified following the semi-quantitative method of Biscaye [19]; total organic carbon (TOC) was determined following the wet oxidation method [20]; Magnetic susceptibility measurements of sediments were carried out and various parameters were calculated in terms of magnetic concentration, mineralogy and grain size as summarized by Thomson and Oldfield [21] and Oldfield [22]; total/bulk metal concentrations in sediments, sediment samples were digested following the procedure given by [23] and analyzed using an Atomic Absorption spectrophotometer (Flame AAS). Sediment standard reference 2702 was used to compute percentage recovery. Further, analysis of speciation of metals was carried out following the procedure developed by Tessier et al. [24] involving sequential chemical extractions of sediments. Five fractions namely exchangeable, bound to carbonates, bound to Fe-Mn oxides, bound to organic matter/ sulfides, and residual was employed. Metals in sediment associated organisms and mangrove pneumatophores collected from the specified sampling locations were also studied. Al, Fe, Mn, Cr, Co, Ni, Cu and Zn were studied for total sediment, speciation and bioaccumulation. Statistical analysis and sediment quality assessment were applied to understand level of metal concentration. Selected sediment cores were dated using lead dating method.

Observations and Discussion

Our investigations on the estuaries along central west coast of India with special references to sediment components and metal content in the sediment with an aim to understand the depositional environment in estuarine mangrove and mudflat sediments along with source, behavior, contamination and bioavailability of metals [25-41] have provided significant results. Also, an attempt was made to study on bioaccumulation and bioremediation potential in an estuary [42,43]. Further, analyses have also been carried out using different proxies namely diatoms, sediment grain size, isotopes to understand the recent past climate variations [44,45].

From our studies along central west coast of India it was clear that




Figure 2: Classic estuarine zonation depicted from the head region where fluvial processes dominate, to the mid- and mouth regions where tidal and wave processes are dominant controlling physical forces, respectively. Differences in the intensities and sources of physical forcing throughout the estuary also result in the formation of distinct sediment facies [1,13].

there were considerable variations in the depositional environment with time. The geomorphology of the estuaries, rainfall, river runoff, construction of dams, bridges and other anthropogenic activities have considerably influenced the depositional environment. It is prominently noted that there is a strong transition in the estuarine environment from fresh water dominated in the past to marine inundated in recent times. These changes were mainly attributed to the rise in sea level, rainfall pattern, decrease in runoff due to the construction of dams in recent times etc. Several estuaries such as Vaitarna, Rajapur, Mandovi, Sharavathi are dammed in the upstream regions for diversion of river water and irrigational purposes in the recent years that resulted in reduced fresh water runoff. This in turn enhanced the tidal surge into the estuary and hence more saline water intrusion towards the upper reaches of the estuaries in recent times. Thus, the mixing processes were affected leading to increased flocculation and deposition of fine grained sediments in the recent years. Further, anthropogenic activities in the catchment areas of the estuaries have considerably affected sedimentation. Activities like mining and transportation of ores add considerable amount of material in to the estuaries.

The characteristics of an estuary are determined by the dynamics of various processes and the sediment sources. Material is imported from the river and its catchments, and sea into the estuary where the transformation of material takes place. Post transformation, a part of the material such as a particulate matter is retained in the estuary in sediments whereas the dissolved material is exported to the sea [46]. The main sediment sources of an estuary are from existing base material, terrigenous material held in the catchment and sand transported from the open-coast marine environment. In addition particulate and dissolved matter composed of organic and inorganic material is added in to the estuary that may be supplied naturally or from anthropogenic sources. The distribution of the sediments within an estuary is regulated by interactions between the available sediments, bottom morphology and flow hydrodynamics [47]. The dynamics of sediment transport depend on the water circulation, salinity, biological interaction, and sediment type [48]. The interaction among cohesive sediments (mud) is different from that of non-cohesive sediments (sand). Cohesive sediments may aggregate, forming flocs by the flocculation process caused by chemical or biological interaction. Flocculation increases the settling velocity of sediment particles. Chemical flocculation is started by salinity, ions that attach to the small mud particles, cause electronic forces between

the particles, which start aggregating and thus forming a larger mud floc. In contrast, “Biological flocculation” is caused by bacteria and plankton that produce exopolymer and binds mud particles leading to formation of extremely large flocs of ~1000 μm in size [49].

Estuaries contain many different habitats such as shallow open waters, sandy beaches, salt marshes mud and sand flats, rocky shores, mangrove forests, seagrass beds, river deltas and tidal pools. Out of these, mangrove and mudflats have received considerable attention as they are very effective in coastal protection, respond to sea level changes and offer an important habitat for wildlife, food and recreation. Mangroves are a diverse group of trees, palms, shrubs, vines and ferns that have the ability to thrive in waterlogged saline soils that are subjected to regular flooding by tides. Mangrove forests are typically found in the tropical and sub-tropical latitudes, lying between the land and the sea in sheltered coastal areas that are subjected to tidal influence and are among the most productive ecosystems [50,51]. The mangrove ecosystems play an important role in carbon, nitrogen, phosphorus, and sulfur cycles, in addition to providing protection to coastal areas from waves and storms [52]. Mangroves are known for being sites for sediment deposition and are associated carbon and nutrients [53,54]. The mangroves trap sediment by their complex aerial root structure and are an important sink for suspended sediments and thus function as land builders [42,55-59].

Mudflats are coastal wetlands formed in sheltered shores where greater amounts of sediments, detritus are deposited by the rivers or tides. They are frequently associated with estuaries, and are usually situated adjacent to mangroves and comprise around 7 % of total coastal shelf areas [60]. Intertidal mudflats play a critical role in the estuarine exchange of marine and continental supplies of nutrients and sediments. The sediments consist mainly of fine particles, mostly in the silt and clay fraction. Little oxygen penetrates through the cohesive sediments, and an anoxic layer is often present within millimeters of the sediment surface. Mudflats generally support very little vegetation other than green algae. Their biodiversity centers on the range of invertebrates living in the sediment which are biologically productive. The intertidal mudflats support communities characterized by polychaetes, bivalves and oligochaetes and large numbers of birds and fish. Mudflats provide an important nursery and feeding ground for many fish species.

Sediments that are transported by the estuarine waters typically

cover a range of sizes from less than 0.002mm to more than 4mm, with the finer sizes dominant in most estuaries. Estuarine sand is typically composed of quartz, although other minerals such as feldspar or various heavy minerals such as magnetite may be present depending on the sediment source. The fine sediments in estuaries are mixtures of inorganic minerals, organic materials, and biochemicals. Mineral grains usually consist of clays such as montmorillonite, illite, and kaolinite and chlorite, and non-clay minerals like quartz and carbonate. Organic materials comprise of biogenic detritus and microorganisms [61].

The estuarine system is mainly an area of deposition and acts as an important sink for metals in the environment. Metals are supplied to the estuary by natural factors such as chemical leaching of bedrocks, water drainage basins, and runoff from banks while the discharge of urban and industrial waste water, combustion of fossil fuels, mining and smelting operations, waste disposal and transportation activities are the important anthropogenic sources of the metal pollutants [25,27,28,30,34,37,62-66]. Metals brought into the estuary are transferred from solution to sediments by adsorption onto suspended particulate matter, and are deposited with relatively short lag times and tend to get trapped and accumulate in the sediments [67].

The distribution and accumulation of metals are influenced by the sediment texture, mineralogical components and physical transport [68,69]. The metals get assimilated in the sediment along with organic matter, Fe/Mn oxides, sulphide, and clay and thus undergo alterations in their speciation due to geochemical modifications by processes such as dissolution, precipitation, sorption and complexation when discharged into the estuary and form several reactive components [70]. The sediment characteristics such as pH, cation exchange capacity, organic matter content, redox conditions, chloride content and salinity determine metal sorption and precipitation processes, which are associated to the metal mobility, bioavailability and potential toxicity [71]. The organic matter content in the sediments leads to relatively higher metal accumulation [72]. In addition, sediment grain size substantially influences the metal concentration in the estuarine sediments as the clay fractions that have a high specific surface area, favor adsorption processes [73]. Following deposition and burial, metals become subject to a variety of physical, chemical and biological processes which may mix and remobilize the metals into the water column [74] or may be immobilized in the sediments for long periods of time and undergo compaction and diagenesis.

The chemical speciation of metals in sediments involves the identification and quantification of the different forms or phases of the metal present in the sediments and provides advanced information on the potential availability of metals to biota under various environmental conditions [75-78]. The fractionation procedure can indicate the propensity for metals to be remobilized and can help distinguish those metals having a lithogenic origin from those with an anthropogenic origin [79,80]. Tessier et al. [24] devised a fractionation procedure which defined the desired partitioning of trace metals into fractions that are likely to be affected by various environmental conditions i.e. exchangeable, bound to carbonates, bound to iron and manganese oxides, organic matter/sulfide bound and the residual fraction. The metals in the exchangeable fraction are likely to be affected by sorption-desorption processes such as weakly bound to clays, hydrated oxides of iron and manganese and humic acids while the metals in the carbonate fractions can be associated with sediment carbonates and this fraction is susceptible to changes of pH. Together the exchangeable and carbonate fractions are known as the

labile fraction [81]. The third fraction of sediment consists of metals bound to iron and manganese oxides and these oxides are excellent scavengers for trace metals [82] and are thermodynamically unstable under anoxic conditions. The fourth fraction consists of trace metals bound to various forms of organic matter such as detritus, humic and fulvic acids etc, through complexation and peptization phenomenon. A large amount of sulfides are also leached into this fraction. Under oxidizing conditions in natural waters, the degradation of organic matter leads to the release of soluble trace metals. The residual fraction of the sediments consists of primary and secondary minerals which may retain trace metals within their crystal structure and are not released easily into solution. The first four fractions are known as the bioavailable fraction as they exhibit mobility and are potentially available for uptake by organisms. The mobile fractions introduced by anthropogenic activities remain bound to the exchangeable, the carbonate and the easily reducible phases [83]. The sediment-associated metals have the potential to be ecotoxic due to their mobility and bioavailability, and this in turn affects both ecosystems and life through a process of bioaccumulation and biomagnification, respectively [12,68]. Thus, evaluating metal speciation can provide detailed information about the origin, mobilization, contamination risks, biological availability and toxicity of metals [84].

As sediments are often the final repository of metals, accumulation of high concentrations of metals can present a risk to organisms [85]. Metals normally occurring in nature are not harmful to the environment, because they play an essential role in tissue metabolism and growth of plants and animals [86]. However, metals like Cu, Zn, Fe, Co, Mo, Ni, Si, and Sn become predominantly toxic when their level exceeds the limit, and V, Cd, Pb, and Hg are prominently classified as toxic because of their detrimental effect even at low concentrations [87]. Marine organisms can accumulate metals in their tissues that may threaten the health of organisms higher in the food chain through trophic transfer to terrestrial, estuarine and eventually coastal species that become prey for oceanic predators and humans [88,89]. Metals are ingested of metal-enriched sediment and suspended particles during feeding, and by uptake from solution [90]. The ecological risk posed by metal-contaminated sediments depends strongly on the sediment characteristics, specific chemical forms of the metals, influencing their availability to aquatic organisms (bioavailability) and the ability of these organisms to accumulate (bioaccumulation) or remove metals. The efficiency of bioaccumulation via sediment ingestion is dependent on geochemical characteristics of the sediment. Various accumulation patterns have been described regulating the uptake of metals defined by the balance between uptake and excretion rates [91,92] and by detoxification processes usually involving proteins such as metallothioneins [93-95].

Conclusion

Study carried out on estuarine mudflats and mangroves sedimentary sub-environments along central west coast of India revealed that geomorphology, rainfall, river runoff, construction of dams, bridges and other anthropogenic activities have considerably influenced the depositional environment. The cohesive sediments present in these environments protect the coast from erosion and the sediment column can be used to understand past depositional environments, sea level changes, changes in land use-land cover in the catchment area of the rivers and bioavailability of metals in addition to depositional environments.

References

- Bianchi TS. Biogeochemistry of Estuaries. Oxford University Press, Oxford, 2006; 720pp.
- Buddemeier RW, Smith SV, Swaney DP, Crossland CJ, Maxwell BA. Coastal typology: An integrative "neutral" technique for coastal zone characterization and analysis. *Estuarine, Coastal and Shelf Science*. 2008; 77: 197-205.
- American Geological Institute. Glossary of Geology and Related Sciences with Supplement. Am. Geol. Inst., Washington, D.C., 2nd Ed. 1960; 371pp.
- Cameron WM, Pritchard DW. *Estuaries, The Sea*. 1963; 2: 306-324.
- Pritchard DW. What is an estuary: physical viewpoint. *American Association for the Advancement of Science*. 1967; 83: 3-5.
- Yu X, Yan Y, Wang WX. The distribution and speciation of trace metals in surface sediments from the Pearl River Estuary and the Daya Bay, Southern China. *Marine Pollution Bulletin*. 2010; 60: 1364-1371.
- Liu WX, Li XD, Shen ZG, Wang DC, Wai OWH, Li YS. Multivariate statistical study of heavy metal enrichment in sediments of the Pearl River Estuary. *Environmental Pollution*. 2003; 121: 377-388.
- Pritchard DW. Estuarine hydrography. In: H.E. Landsberg (Editor), *Advances in Geophysics*. Academic Press, New York. 1952; 1: 243-280.
- Valle-Levinson A. Contemporary issues in estuarine physics. Cambridge University Press, 2010.
- Hayes MO. Morphology of sand accumulation in estuaries. In: L.E. Cronin (Editor), *Estuarine research*, Vol. 2, Academic press, New York. 1975; 3-22.
- Fairbridge RW. The Estuary: its definition and geodynamic cycle. In: Olausson, E., Cato, I. (Eds.), *Chemistry and Biogeochemistry of Estuaries*. Wiley, New York. 1980; 1-35.
- Ip, CC, Li XD, Zhang G, Wai OW, Li YS. Trace metal distribution in sediments of the Pearl River Estuary and the surrounding coastal area, South China. *Environmental Pollution*. 2007; 147: 311-323.
- Dalrymple RW, Zaitlin BA, Boyd R. A conceptual model of estuarine sedimentation. *Journal of Sedimentary Petrology*. 1992; 62: p.116.
- Zingde MD, Ram A, Sharma P, Abidi SAH. Seawater intrusion and behaviour of dissolved boron, fluoride, calcium, magnesium and nutrients in Vashisti estuary. In: Agarwal VP, Sharma CB, Abidi SAH and Zingde MD (eds.), *Complex carbohydrates & advances in biosciences*. 1995; 563-580.
- Wensink H. Newer paleomagnetic results of the Deccan Traps, India. *Tectonophysics*. 1973; 17: 41-59.
- Naqvi SM. Geology and evolution of the Indian Plate (from Hadean to Holocene - 4 Ga to 4 Ka). Capital publishing company, New Delhi. 2005; 450.
- Folk RL. In *Petrology of sedimentary rocks*. Austin, Texas: Hemphill. 1974; 182pp.
- Rao VP, Rao BR. Provenance and distribution of clay minerals in the sediments of the western continental shelf and slope of India. *Continental Shelf Research*. 1995; 15: 1757-1771.
- Biscaye PE. Mineralogy and sedimentation of recent deep sea clay in the Atlantic Ocean and adjacent seas and oceans. *Geological Society of America Bulletin*. 1965; 76: 803-832.
- Walkley A, Black IA. An examination of the degtjareff method for the determining organic matter and a proposed modification of the chromic acid titration method. *Soil Science*. 1934; 37: 29-38.
- Thomson R, Oldfield F. *Environmental magnetism*. Allen and Unwin, London. 1986; pp. 227.
- Oldfield F. *Environment magnetism: The range of applications In Walden*, J., Smith, J.P. and Oldfield, F. (eds.) *Environmental magnetism, a practical guide*, Quaternary research association, Technical Guide. 1999; 6: 212-222.
- Jarvis IJ, Jarvis K. Rare earth element geochemistry of standard sediments: a study using inductively coupled plasma spectrometry. *Chemical Geology*. 1985; 53: 335-344.
- Tessier A, Campbell PG, Bisson M. Sequential extraction procedure for the speciation of particulate trace metals. *Analytical chemistry*. 1979; 51: 844-851.
- Nayak GN, Noronha-D'Mello CA, Pande A, Volvoikar SP. Understanding sedimentary depositional environments through geochemical signatures of a Tropical (Vaghotan) estuary, West Coast of India. *Environmental Earth Sciences*. 2016; 75: 1-15.
- Fernandes MC, Nayak GN. Speciation of metals and their distribution in tropical estuarine mudflat sediments, southwest coast of India. *Ecotoxicology and environmental safety*. 2015; 122: 68-75.
- Fernandes MC, Nayak GN, Pande A, Volvoikar SP, Dessai DRG. Depositional environment of mudflats and mangroves and bioavailability of selected metals within mudflats in a tropical estuary. *Environmental Earth Sciences*. 2014; 72: 1861-1875.
- Noronha-D'Mello CA, Nayak GN. Geochemical characterization of mangrove sediments of the Zuari estuarine system, West coast of India. *Estuarine, Coastal and Shelf Science*. 2015; 167: 313-325.
- Nasnodkar MR, Nayak GN. Processes and factors regulating the distribution of metals in mudflat sedimentary environment within tropical estuaries, India. *Arabian Journal of Geosciences*. 2015; 8: 9389-9405.
- Pande A, Nayak GN. Depositional environment and elemental distribution with time in mudflats of Dharamtar creek, west coast of India. *Indian Journal of Geo-Marine Sciences*. 2013b; 42: 360-369.
- Siraswar R, Nayak GN. Mudflats in lower middle estuary as a favorable location for concentration of metals, west coast of India. *Indian Journal of Marine Sciences*. 2011; 40: 372.
- Singh KT, Nayak GN. Sedimentary and geochemical signatures of depositional environment of sediments in mudflats from a microtidal Kalinadi estuary, central west coast of India. *Journal of Coastal Research*. 2009; 641-650.
- Fernandes MC, Nayak GN. Role of sediment size in the distribution and abundance of metals in a tropical (Sharavati) estuary, west coast of India. *Arabian Journal of Geosciences*. 2016; 9: 1-13.
- Fernandes LL, Nayak GN. Characterizing metal levels and their speciation in intertidal sediments along Mumbai coast, India. *Marine pollution bulletin*. 2014; 79: 371-378.
- Fernandes LL, Nayak GN. Heavy metals contamination in mudflat and mangrove sediments (Mumbai, India). *Chemistry and Ecology*. 2012; 28: 435-455.
- Fernandes LL, Nayak GN. Sources and factors controlling the distribution of metals in mudflat sedimentary environment, Ulhas Estuary, Mumbai. *Indian Association of Sedimentologists*. 2010; 29: 71-83.
- Volvoikar SP, Nayak GN. Impact of industrial effluents on geochemical association of metals within intertidal sediments of a creek. *Marine pollution bulletin*. 2015; 99: 94-103.
- Volvoikar SP, Nayak GN. Factors controlling the distribution of metals in intertidal mudflat sediments of Vaitarna estuary, North Maharashtra coast, India. *Arabian Journal of Geosciences*. 2014b; 7: 5221-5237.
- Volvoikar SP, Nayak GN. Evaluation of impact of industrial effluents on intertidal sediments of a creek. *International Journal of Environmental Science and Technology*. 2013; 10: 941-954.
- Singh KT, Nayak GN, Fernandes LL, Borole DV, Basavaiah N. Changing environmental conditions in recent past—Reading through the study of geochemical characteristics, magnetic parameters and sedimentation rate of

- mudflats, central west coast of India. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 2014; 397: 61-74.
41. Singh KT, Nayak GN, Fernandes LL. Geochemical evidence of anthropogenic impacts in sediment cores from mudflats of a tropical estuary, Central west coast of India. *Soil and Sediment Contamination*. 2013; 22: 256-272.
42. Noronha-D'Mello CA, Nayak GN. Assessment of metal enrichment and their bioavailability in sediment and bioaccumulation by mangrove plant pneumatophores in a tropical (Zuari) estuary, west coast of India. *Marine Pollution Bulletin*. 2016; 110: 221-230.
43. Dias HQ, Nayak GN. Geochemistry and bioavailability of mudflats and mangrove sediments and their effect on bioaccumulation in selected organisms within a tropical (Zuari) estuary, Goa, India. *Marine pollution bulletin*. 2016; 105: 227-236.
44. Pande A, Nayak GN, Prasad V, PrakashBabu C. Geochemical and diatom records of recent changes in depositional environment of a tropical wetland, central west coast of India. *Environmental Earth Sciences*. 2015; 73: 5447-5461.
45. Volvoikar SP, Nayak GN, Mazumdar A, Peketi A. Reconstruction of depositional environment of a tropical estuary and response of $\delta^{13}C$ org and TOC/TN signatures to changing environmental conditions. *Estuarine, Coastal and Shelf Science*. 2014; 139: 137-147.
46. Turner S, Riddle B. Estuarine Sedimentation and Vegetation – Management Issues and Monitoring Priorities. *Environment Waikato Internal Series 2001/05. Document #: 686944*. 2001.
47. Frey RW, Howard JD. Mesotidal estuarine sequences: a perspective from the Georgia Bight. *Journal of Sedimentary Research*. 1986; 56.
48. Wang XH, Andutta FP. Sediment transport dynamics in ports, estuaries and other coastal environments. *Intech. Open Access Publisher*. 2013.
49. Wolanski E, Andutta F, Delhez E. Estuarine hydrology. In *Encyclopedia of Lakes and Reservoirs*, Springer Netherlands. 2012; 238-249.
50. Kathiresan K, Bingham BL. Biology of mangroves and mangrove ecosystems. *Advances in marine biology*. 2001; 40: 81-251.
51. Kathiresan K. Greening the blue mud. *Revista de biología tropical*. 2002; 50: 869-874.
52. Meng X, Xia P, Li Z, Liu L. Mangrove forest degradation indicated by mangrove-derived organic matter in the Qinzhou Bay, Guangxi, China, and its response to the Asian monsoon during the Holocene climatic optimum. *Acta Oceanologica Sinica*. 2016; 35: 95-100.
53. Eyre B. Nutrients in the sediments of a tropical north-eastern Australian estuary, catchment and nearshore coastal zone. *Marine and Freshwater Research*. 1993; 44: 845-866.
54. Furukawa K, Wolanski E. Sedimentation in mangrove forests. *Mangroves and salt marshes*. 1996; 1: 3-10.
55. Woodroffe C. Mangrove sediments and geomorphology. In Robertson, A. I. and Alongi, D.M. (eds.), *Tropical mangrove ecosystem*, American Geophysical Union, Washington. 1992; 7-41.
56. Wolanski E, Mazda Y, Ridd P. Mangrove hydrodynamics. In: Robertson, A.I. and Alongi, D.M., (eds.). *Tropical mangrove ecosystem*, American Geophysical Union, Washington D.C. 1992; 436-462.
57. Wolanski E. Physical oceanographic processes of the Great Barrier Reef. CRC Press. 1994.
58. Wolanski E. Transport of sediment in mangrove swamps. *Hydrobiologia*. 1995; 295: 31-42.
59. Furukawa K, Wolanski E, Mueller H. Currents and sediment transport in mangrove forests. *Estuarine, Coastal and Shelf Science*. 1997; 44: 301-310.
60. Stutz ML, Pilkey OH. Global distribution and morphology of deltaic barrier island systems. *Journal of Coastal Research*. 2002; 36: 694-707.
61. McNally WH, Mehta AJ. Sediment Transport and Deposition in Estuaries (Sample Chapter). In *Encyclopedia of Life Support Systems (EOLSS): Coastal Zones and Estuaries*. 2004.
62. Abdullah AR, Tahir NM, Loong TS, Hoque TM, Sulaiman AH. The GEF/UNDP/IMO Malacca Straits demonstration project: sources of pollution. *Marine Pollution Bulletin*. 1999; 39: 229-233.
63. Shazili NAM, Yunus K, Ahmad AS, Abdullah N, Rashid MKA. Heavy metal pollution status in the Malaysian aquatic environment. *Aquatic Ecosystem Health and Management*. 2006; 9: 137-145.
64. Dragun Z, Roje V, Mikac N, Raspor B. Preliminary assessment of total dissolved trace metal concentrations in Sava River water. *Environmental Monitoring and Assessment*. 2009; 159: 99-110.
65. Pardo R, Barrado E, Lourdes P, Vega M. Determination and speciation of heavy metals in sediments of the Pisuerga River. *Water Research*. 1990; 24: 373-379.
66. Zhou J, Ma D, Pan J, Nie W, Wu K. Application of multivariate statistical approach to identify heavy metal sources in sediment and waters: a case study in Yangzhong, China. *Environmental Geology*. 2008; 54: 373-380.
67. Spencer KL, Cundy AB, Croudace IW. Heavy metal distribution and early-diagenesis in salt marsh sediments from the Medway Estuary, Kent, UK. *Estuarine, Coastal and Shelf Science*. 2003; 57: 43-54.
68. Buccolieri A, Buccolieri G, Cardellicchio N, Dell'Atti A, Di Leo A, Maci A. Heavy metals in marine sediments of Taranto Gulf (Ionian Sea, southern Italy). *Marine chemistry*. 2006; 99: 227-235.
69. Marchand C, Lallier-Verges E, Baltzer F, Albéric P, Cossa D, Baillif P. Heavy metals distribution in mangrove sediments along the mobile coastline of French Guiana. *Marine chemistry*. 2006; 98: 1-17.
70. Lim WY, Aris AZ, Zakaria MP. Spatial variability of metals in surface water and sediment in the Langat river and geochemical factors that influence their water-sediment interactions. *The Scientific World Journal*. 2012.
71. Du Laing G, Bogaert N, Tack FM, Verloo MG, Hendrickx F. Heavy metal contents (Cd, Cu, Zn) in spiders (*Pirata piraticus*) living in intertidal sediments of the river Scheldt estuary (Belgium) as affected by substrate characteristics. *Science of the Total Environment*. 2002; 289: 71-81.
72. Zhong AP, Guo SH, Li FM, Gang LI, Jiang KX. Impact of anions on the heavy metals release from marine sediments. *Journal of Environmental Sciences*. 2006; 18: 1216-1220.
73. Thuy HTT, Tobschall HJ, An PV. Trace element distributions in aquatic sediments of Danang–Hoian area, Vietnam. *Environmental Geology*. 2000; 39: 733-740.
74. Lee SV, Cundy AB. Heavy metal contamination and mixing processes in sediments from the Humber Estuary, Eastern England. *Estuarine, Coastal and Shelf Science*. 2001; 53: 619-636.
75. Álvarez-Iglesias P, Rubio B. Redox status and heavy metal risk in intertidal sediments in NW Spain as inferred from the degrees of pyritization of iron and trace elements. *Marine pollution bulletin*. 2009; 58: 542-551.
76. Fytianos K, Lourantou A. Speciation of elements in sediment samples collected at lakes Volvi and Koronia, N. Greece. *Environment International*. 2004; 30: 11-17.
77. Gao X, Chen S, Long A. Chemical speciation of 12 metals in surface sediments from the northern South China Sea under natural grain size. *Marine Pollution Bulletin*. 2008; 56: 770-797.
78. Rauret G. Extraction procedures for the determination of heavy metals in contaminated soil and sediment. *Talanta*. 1998; 46: 449-455.
79. Förstner U, Ahlf W, Calmano W, Kersten M, Schoer J. Assessment of mobility in sludges and solid wastes. In: Broecker, J.A.C., Gucer, S. and Adams, F. (eds.), *Metal speciation in the environment*. NATO ASI Series G, Ecological Sciences, 23, Springer, Berlin. 1990; 1-41.

80. Korfali SI, Jurdi MS. Speciation of metals in bed sediments and water of Qaraaoun Reservoir, Lebanon. *Environmental monitoring and assessment*. 2011; 178: 563-579.
81. Perin G, Craboledda L, Lucchese M, Cirillo R, Dotta L, Zanetta ML, et al. Heavy metal speciation in the sediments of Northern Adriatic Sea. A new approach for environmental toxicity determination. In: Lakkas, T.D., (ed). *Heavy Metals in the Environment*, CEP Consultants Edinburg. 1985; 454-456.
82. Gutierrez M. Trace element concentration patterns in sediments of the lower Rio Conchos, Mexico. *Water, air and soil pollution*. 2000; 121: 259-270.
83. Nair CK. Chemical partitioning of trace metals in sediments of a tropical estuary. Thesis submitted to The Cochin University of Science and Technology. 1992; pp 4.
84. Yang J, Cao L, Wang J, Liu C, Huang C, Cai W, et al. Speciation of metals and assessment of contamination in surface sediments from Daya Bay, South China Sea. *Sustainability*. 2014; 6: 9096-9113.
85. Casado-Martinez MC, Smith BD, Rainbow PS. Assessing metal bioaccumulation from estuarine sediments: comparative experimental results for the polychaete *Arenicola marina*. *Journal of Soils and Sediments*. 2013; 13: 429-440.
86. Amundsen PA, Staldivik FJ, Lukin AA, Kashulin NA, Popova OA, Reshetnikov YS. Heavy metal contamination in freshwater fish from the border region between Norway and Russia. *Science of the Total Environment*. 1997; 201: 211-224.
87. Michael HC. In: *Encyclopedia of Earth*. Monosson Cleveland C, editor. Washington DC: National Council for Science and the Environment. Heavy metal. 2010.
88. Boyle D, Brix KV, Amlund H, Lundebye AK, Hogstrand C, Bury NR. Natural arsenic contaminated diets perturb reproduction in fish. *Environmental science and technology*. 2008; 42: 5354-5360.
89. Cheung MS, Wang WX. Analyzing biomagnification of metals in different marine food webs using nitrogen isotopes. *Marine Pollution Bulletin*. 2008; 56: 2082-2088.
90. Luoma SN. Bioavailability of trace metals to aquatic organisms—a review. *Science of the total environment*. 1983; 28: 1-22.
91. Luoma SN, Rainbow PS, Luoma S. *Metal contamination in aquatic environments: science and lateral management*. Cambridge University Press. 2008; pp. 141-150.
92. Rainbow PS. Trace metal concentrations in aquatic invertebrates: why and so what? *Environmental Pollution*. 2002; 120: 497-507.
93. Casado-Martinez MC, Smith BD, Luoma SN, Rainbow PS. Metal toxicity in a sediment-dwelling polychaete: threshold body concentrations or overwhelming accumulation rates? *Environmental Pollution*. 2010; 158: 3071-3076.
94. Greim H, Snyder R. *Toxicology and risk assessment: a comprehensive introduction*. John Wiley and Sons, (eds.). 2008; 1-18.
95. Walker CH, Sibly RM, Hopkin SP, Peakall DB. *Principles of ecotoxicology*, CRC press. 2012.