

# SCIENTIFIC REPORTS



OPEN

## Light triggered encapsulation and release of C<sub>60</sub> with a photoswitchable TPE-based supramolecular tweezers


Mousumi Samanta<sup>1</sup>, Anushri Rananaware<sup>2</sup>, Dinesh N. Nadimetta<sup>3</sup>, Sk. Atiur Rahaman<sup>1</sup>, Monochura Saha<sup>1</sup>, Ratan W. Jadhav<sup>3</sup>, Sheshanath V. Bhosale<sup>3</sup> & Subhajit Bandyopadhyay<sup>1</sup>

Stimuli responsive hosts for C<sub>60</sub> can control its binding and release on demand. A photoswitchable TPE based supramolecular host can encapsulate C<sub>60</sub> in the Z-form with a markedly different visual change in the colour. In addition, the Z-1 bound C<sub>60</sub> has been characterized by various spectroscopic methods and mass spectrometry. Upon exposure to visible light (>490 nm), the host switches to the E-form where the structural complementarity with the guest is destroyed as a result of which the C<sub>60</sub> is disassembled from the host. The results described herein reveals an actionable roadmap to pursue further advances in component self-assembly particularly light-induced association and dissociation of a guest molecule.

The discovery of fullerene is marked as an epoch in the field of organic nanomaterials<sup>1–3</sup>. Since C<sub>60</sub> is a spherical  $\pi$  framework devoid of any functional groups, construction of its receptor has been a challenge for chemists. A number of host molecules for C<sub>60</sub> stands as witnesses of ingenuity of molecular architecture<sup>4</sup>. Complementarity of the spherical shape of the buckyball has been achieved by challenging synthesis of molecular bowls, hoops, peapods and several other flexible structures that undergo induced-fit around the molecule. Cycloparaphenylene acetylenes offer supramolecular complexation with fullerene derivatives<sup>5</sup>. Platinum-based molecular cages has also been reported to encapsulate the C<sub>60</sub><sup>6</sup>. Aromatic molecules such as cycloparaphenylene rings encapsulate C<sub>60</sub> as well as other fullerene derivatives<sup>7,8</sup>. An aza-buckybowl based system have been reported to act as efficient receptors for C<sub>60</sub> and C<sub>70</sub> with large association constant<sup>9</sup>. Larger aromatics such as cyclochrysenylenes form molecular “peapods”. The complementarity of the convex surface of the buckyball has been achieved by concave bowl-shaped molecules such as corannulenes, sumanenes and even with an expanded rosin derivative<sup>10–13</sup>. Even a nitrogen-containing buckybowl and its assembly with C<sub>60</sub> has recently been reported<sup>14</sup>. Non-planar hydrocarbons such as triptycene also form complexes with fullerenes due to geometrical complementarity of their concave shapes to the spherical surface of the C<sub>60</sub><sup>15</sup>. Fusion of more than one receptor units in a single host molecule displayed enhanced affinity towards the fullerene guest. For instance, a receptor bearing two corannulene units has been reported as a “bucky-catcher” with a high binding constant<sup>16</sup>.

Stimuli-responsive supramolecular host for fullerenes can open up the possibilities for handling the bucky balls and other carbon-based frameworks with superior controls using the stimuli such as pH, electrical or magnetic fields, electrochemical and photonic signals<sup>17–21</sup>. Light is one of the most extensively used stimuli because of its non-invasive nature and easy regulation by the precise focus of its exposure area, and adjustment of its wavelength and intensity<sup>22,23</sup>. Several light-triggered and redox-triggered release of guest molecules have been reported in recent years<sup>24–27</sup>. The selective recognition of C<sub>60</sub> by a bispyridine ligand with embedded anthracene panels in presence of Ag(I) ion and its light-mediated subsequent release was studied<sup>28</sup>. Reversible host-guest complexation and decomplexation triggered by light can be achieved by the incorporation of photochromic units in the supramolecular systems<sup>29</sup>. Azobenzene is a robust photoresponsive molecule which can exhibit significant structural and chemical modification upon exposure of UV and visible light. These photoswitches have been widely used for the rapid and exact modulation of several biological processes<sup>30–45</sup>.

<sup>1</sup>Indian Institute of Science Education and Research (IISER) Kolkata, Mohanpur, Nadia, WB, 741246, India. <sup>2</sup>School of Science, RMIT University, Melbourne, Victoria, 3001, Australia. <sup>3</sup>School of Chemical Sciences, Goa University, Taleigao Plateau, Goa, 403206, India. Mousumi Samanta, Anushri Rananaware and Dinesh N. Nadimetta contributed equally. Correspondence and requests for materials should be addressed to S.V.B. (email: [svbhosale@unigoa.ac.in](mailto:svbhosale@unigoa.ac.in)) or S.B. (email: [sb1@iiserkol.ac.in](mailto:sb1@iiserkol.ac.in))



**Figure 1.** The chemical structure: The structure of the molecule **1** and the crystal structure (50% probability ellipsoids) (CCDC Number 1839412).

A theoretical study of host-guest interactions between  $C_{60}$  and a photoresponsive group containing nano-rings host was investigated and further experimental synthesis of photoresponsive hosts have been predicted<sup>46</sup>. However, reversible binding and release of  $C_{60}$  still remain a challenge.  $C_{60}$  is known to form stable supramolecular complexes with several compounds having flexible phenyl rings<sup>47</sup>. Crystal structures showing  $C_{60}$ -tetraphenylethene (TPE) bears the witness of the non-covalent van der Waals interactions between the two species<sup>48</sup>. Recent years, TPE have been widely used due to its abnormal properties i.e. non-luminescent in a solution state and highly emissive upon aggregation and solid state — so called aggregation induced emission (AIE) active chromophore<sup>49,50</sup>.


## Results

Incorporating a photoswitchable azobenzene unit between two TPE moieties, we have synthesized the azobenzene-TPE (**1**, Fig. 1) that contains the elements of supramolecular interactions to bind to  $C_{60}$ . Binding through supramolecular association is much coveted because it does not perturb the electronic structure of the guest significantly. The azobenzene-TPE photoswitch offers the control of the geometry of the receptor allowed reversible supramolecular interaction with  $C_{60}$ .


**Synthesis.** Azobenzene-TPE **1** was prepared by reacting 1,2-bis(4-bromophenyl)diazene with 3 equiv. of (4-(1,2,2-triphenylvinyl)phenyl)boronic acid in degassed 1,2-dimethoxyethane/2 M  $Na_2CO_3$  (3:1, v/v) using  $Pd(PPh_3)_4$  as a catalyst at 100 °C for 24 h yielded **1** in 69.5% (for details see Supplementary Information).

**Photoisomerization studies of 1.** The molecule **1** exhibit *E-Z* photochromism where the forward and the backward reactions are fully reversible upon exposure to UV and visible light respectively. Upon irradiation with 254 nm light, the  $n - \pi^*$  band of the *E* isomer can be selectively excited leading to the conversion to *Z* isomer as shown in Fig. 2a. This phenomenon can be visualized by the naked eye (Fig. 2b). Under the exposure of 254 nm UV light the band at 400 nm having a molar extinction coefficient ( $\epsilon$ ) of  $\sim 1.04 \times 10^5 M^{-1}cm^{-1}$  becomes broad and the intensity of the band increases with the exposure time. A new broad peak at 550 nm ( $\epsilon = \sim 9.8 \times 10^3 M^{-1}cm^{-1}$ ) arises due to the  $n - \pi^*$  transition of the *Z* isomer which increases in intensity with the 254 nm radiation. The photoisomerization reaction was accompanied by a change in colour that was visually noticeable. The yellow colour of the *E* isomer was transformed to intense orange under isomerization to the *Z* isomer (Fig. 2c) The *E* to *Z* conversion monitored by  $^1H$  NMR spectroscopy followed a first order kinetics with a rate constant of  $6.6 \times 10^{-2} m^{-1}$ . The reverse *Z-E* conversion reaction was achieved with visible light with a 400 nm cut off filter. The rate constant for the conversion obtained with the NMR methods was  $8.3 \times 10^{-4} m^{-1}$ . Furthermore, the fluorescence spectra of *E-1* at  $\lambda_{ex} = 405$  nm displayed a broad band with two peaks at 440 and 470 nm ( $\Phi_F = 0.01$ ) and a shoulder at  $\sim 550$  nm. In the *Z* isomer, the fluorescence intensity was less intense ( $\Phi_F = 0.004$ ) with a single band at 480 nm as shown in Fig. 2d.

**NMR studies for isomerization.** The *E-Z* isomerization of **1** under the exposure of 254 nm light was monitored by  $^1H$  spectroscopy (Fig. 3) using  $CS_2$  as the solvent with a small amount of  $CDCl_3$  for the purpose of locking the instrument. The characteristic signals of the azobenzene moiety of *E*-isomer of **1** were observed at  $\delta$  7.87 ( $H_a$ ), 7.58 ( $H_b$ ), and 7.32 ( $H_c$ ). Conversion to the *Z*-isomer upon exposure to 254 nm UV light triggered an upshifted field shift of the  $^1H$  resonances and to  $\delta$  values (a) and (b) protons appear at 7.34 ( $H_c'$ ), 7.19 ( $H_b'$ ) and




**Figure 2.** Photoisomerization studies of **1**: (a) chemical illustration of *E-Z* isomerization of the molecule **1**. (b) Naked eye visualisation of *E-Z* isomers. (c) The changes in absorption spectra of the *E* to *Z* isomerization of the molecule **1** (6  $\mu\text{M}$ ) in  $\text{CS}_2$ . (d) Fluorescence spectra of the azobenzene-TPE **1** in both *E* and *Z* forms.



**Figure 3.**  $^1\text{H}$  NMR spectra of *E* to *Z* isomerisation of the molecule **1**.


6.83 ( $\text{H}_a'$ ). The TPE protons appeared as multiplets and their change in the  $^1\text{H}$  NMR spectra was not identifying because of the presence of overlapping multiplets. The composition under the irradiation in the NMR tube corresponded to a ca. 3:1 ratio of the *Z*:*E* isomer.



**Figure 4.** Encapsulation and release of  $C_{60}$  by the Z form of the molecule. (a) structural illustration of host-guest complex and release of guest. (b,c) Absorbance spectra of the molecule **1** ( $6\ \mu\text{M}$ ) in Z forms with  $C_{60}$  and E form with  $C_{60}$  ( $6\ \mu\text{M}$ ) in  $\text{CS}_2$ , respectively. (d,e) Fluorescence spectra of the molecule **1** ( $6\ \mu\text{M}$ ) in Z forms with  $C_{60}$ . (f) E form with  $C_{60}$  (0 to  $24\ \mu\text{M}$ ) in  $\text{CS}_2$ , respectively. (g) Changes in  $^{13}\text{C}$  NMR spectra of  $C_{60}$  in presence and in absence of Z-isomer of the molecule **1**. (h) The MALDI-TOF MS of a sample containing Z-1. $C_{60}$  indicated the presence of the species perfectly matches well with the calculated isotopic pattern. Changes in (i) UV-vis spectra and (j) fluorescence spectra of Z form of the molecule **1** with  $C_{60}$  under the exposure of visible light in  $\text{CS}_2$ , which clearly shows release of  $C_{60}$  from Z form i.e. converting Z to E isomer with excitation at  $\lambda_{\text{max}} > 490$  nm.

**Light triggered encapsulation of  $C_{60}$  through Photoisomerization.** Upon addition of  $C_{60}$  to Z-1, the formation of Z-1. $C_{60}$  was apparent from the instant change in the absorption spectra. Apart from the obvious change in the visual appearance of the solution Z-1 with  $C_{60}$  upon mixing (Fig. 4b), the spectrum shows an increase in the peak at 540 nm in presence of  $C_{60}$ . The increase in absorption band in the region  $>470$  nm indicates the interaction between Z form of the molecule and  $C_{60}$ . The chemical model illustrated in Fig. 4a. In contrary, such changes were not observed with E-1 (Fig. 4c). In fluorescence spectra, (Fig. 4d,e) a new band at 725 nm was observed in the presence of  $C_{60}$ . In addition, the original fluorescence band of Z-1 along with the shoulder at 520 nm was found to decrease in intensity. A plot of continuous variation (Job's plot) monitored at 460 nm clearly revealed an association with 1:1 stoichiometry of the host and the guest  $C_{60}$  (Supplementary Fig. S1). The binding constant of  $C_{60}$  with both Z-1 and for E-1 has been determined from the emission data. As expected the association constant of  $C_{60}$  with Z-1 was found to be  $4.02 \times 10^4\ \text{M}^{-1}$  (Supplementary Fig. S2) and for E-1 with  $C_{60}$  the value was less by an order in magnitude with a binding constant of  $1.76 \times 10^3\ \text{M}^{-1}$ . The interaction between Z-1 and  $C_{60}$  was also clearly visible with  $^{13}\text{C}$  NMR spectroscopy (Fig. 4g). Upon addition of  $C_{60}$  to Z-1 in  $\text{CDCl}_3$  and  $\text{CS}_2$  (1:10, v/v) the peak of the  $C_{60}$  at 143.0 ppm underwent an upfield shift to 142.9 ppm because of the encapsulation within the aromatic rings of Z-1. The change in the  $^{13}\text{C}$  resonance of the host also provided an insight of the binding site within Z-1 and  $C_{60}$  host-guest complex formation, respectively<sup>5,6</sup>. Importantly, The MALDI-TOF MS of a sample containing Z-1. $C_{60}$  indicated the presence of the species that matches well with the expected isotopic pattern (Fig. 4h).

**Light-triggered release of  $C_{60}$  from the Z form of the molecule.** The Z form of the molecule **1** in the presence of  $C_{60}$  was exposed to  $>490$  nm visible light, then the broad characteristic charge transfer band in the region  $>470$  nm progressively diminished upon irradiation Fig. 4i). In fluorescence spectra, the band at  $>700$  nm



**Figure 5.** Molecular modeling studies of the encapsulation of  $C_{60}$  with the Z form of the molecule **1**.

also gradually disappeared (Fig. 4j). The UV-vis spectra indicated near quantitative conversion of the azobenzene unit to the *E* form. This was confirmed by comparison of the spectra (Fig. 4c) with the one obtained upon addition of  $C_{60}$  to the pure *E* form of **1**.  $^1H$  spectroscopic experiments conducted with **1-Z** and  $C_{60}$  also pointed towards the formation of a host-guest complex between the two. The facile release of the encapsulated guest molecule ( $C_{60}$ ) takes place in presence of the visible light. The sharp characteristics changes in NMR spectra also prove the encapsulation and release of  $C_{60}$  by the Z form of the molecule. The  $^1H$  NMR study was performed with the Z form of the molecule **1** in the presence one equivalent of  $C_{60}$  in  $CS_2$ . Although the changes in the  $^1H$  NMR (Supplementary Fig. S3) was less prominent except for the diminished peaks at 7.34, 7.19 and 6.83, the change in the  $^{13}C$  NMR was clear. There the peak for the  $C_{60}$  at 143.0 shifted upfield to 142.9 ppm (Fig. 4g). This change was reversible under the influence of the visible light. This is anticipated since the  $C_{60}$  upon encapsulation by the host TPE groups in Z form experiences a shielding effect which causes an upfield shift to the  $C_{60}$  nuclei. This change was reversible under exposure to visible light which was also observed earlier with absorption spectroscopy as described earlier.

A three dimensional model of the system obtained from computational simulation using DFT calculation at the B3LYP level displayed clear interactions between the receptor and the fullerene. The  $C_{60}$  molecule fits in perfectly within the aromatic cavity of the **Z-1** form. Several phenyl rings of the molecule puckered around to accommodate the  $C_{60}$  with a perfect shape complementarity to form the supramolecular assembly (Fig. 5).

## Discussion

To sum-up, our salient findings of this work are as follows.

1. A TPE-linked azobenzene based photoswitchable molecule has been synthesized and characterized by various spectroscopic techniques.
2. The structural change of the structure upon *E* → *Z* isomerisation of the compound **1** offers the possibility of the formation of a complementary pocket for the accommodation of  $C_{60}$  guest molecule.
3. Host-guest interaction with both the *E* and the *Z* isomers of **1** and the  $C_{60}$  molecule has been investigated by various spectroscopic techniques.
4. It was observed that the **Z-1**. $C_{60}$  association was pronounced compared to the interaction between the **E-1 isomer** and  $C_{60}$ .
5. The **Z-1**. $C_{60}$  association can be reversed by the exposure of the system with >490 nm light that converts the **Z-1 form** to the **E-1 form**, and thereby weakening the host-guest binding.

Thus this work demonstrates the development of the stimuli responsive host system that can be used for light-induced association and dissociation of a  $C_{60}$  molecule<sup>51</sup>.

## References

1. Dresselhaus, M. S., Dresselhaus, G., Eklund, O. C. Science of fullerenes and carbon nanotubes; academic press: San Diego, CA (1996).
2. Schön, J. H., Kloc, C., Batlogg, B. *Science* **293**, 2432–2434 (2001).
3. Schön, J. H., Kloc, C. & Batlogg, B. Superconductivity at 52 K in hole-doped  $C_{60}$ . *Nature* **408**, 549–552 (2000).
4. Haley, M. M. and Tykwinski, R. R. Carbon-Rich Compounds: From Molecules to Materials Wiley-VCH: Weinheim (2006).
5. Kawase, T., Tanaka, K., Fujiwara, N., Darabi, H. R. & Oda, M. Complexation of a carbon nanoring with fullerene. *Angew. Chem. Int. Ed.* **42**, 1624–1628 (2003).
6. Zhang, M. *et al.* Platinum(II)-Based Convex Trigonal-Prismatic Cages via Coordination-Driven Self-Assembly and  $C_{60}$  Encapsulation. *Inorg. Chem.* **56**, 12498–12504 (2017).
7. Iwamoto, T., Watanabe, Y., Sadahiro, T., Haino, T. & Yamago, S. Sizeselective encapsulation of  $C_{60}$  by [10] cycloparaphenylene: formation of the shortest fullerene-peapod. *Angew. Chem. Int. Ed.* **50**, 8342–8344 (2011).
8. Xia, J., Bacon, J. W. & Jasti, R. Gram-scale synthesis and crystal structure of [8]- and [10] CPP, and the solid-state structure of  $C_{60}$ @ [10]CPP. *Chem. Sci.* **3**, 3018–3021 (2012).
9. Takeda, M. *et al.* Azabuckybowl-Based Molecular Tweezers as  $C_{60}$  and  $C_{70}$  Receptors. *J. Am. Chem. Soc.* **140**, 6336–6342 (2018).
10. Ke, X.-S. *et al.* Expanded Rosarin: A Versatile Fullerene ( $C_{60}$ ) Receptor. *J. Am. Chem. Soc.* **139**, 4627–4630 (2017).
11. Dawe, L. N. *et al.* Corannulene and its penta-tert-butyl derivative co-crystallize 1:1 with pristine  $C_{60}$ -fullerene. *Chem. Commun.* **48**, 5563–5565 (2012).
12. Mehta, G., Shah, R. S. & Ravikumar, K. Towards the design of tricyclopenta [def, jk', pqr] triphenylene ('Sumanene'): a 'bowl-shaped' hydrocarbon featuring a structural motif present in CG0 (Buckminsterfullerene). *J. Chem. Soc. Chem. Comm.* **12**, 1006–1008 (1993).

13. Sakurai, H., Daiko, T. & Hirao, T. A. Synthesis of sumanene, a fullerene fragment. *Science* **301**, 1878 (2003).
14. Yokoi, H. *et al.* Nitrogen-embedded bucky bowl and its assembly with C<sub>60</sub>. *Nat. Commun.* **6**, 8215 (2015).
15. Veen, E. M., Postma, P. M., Jonkman, H. T., Spek, A. L. & Feringa, B. L. Solid state organisation of C<sub>60</sub> by inclusion crystallisation with triptycenes. *Chem. Commun.* 1709–1710 (1999).
16. Sygula, A., Fronczek, F. R., Sygula, R., Rabideau, P. W. & Olmstead, M. M. A double concave hydrocarbon buckycatcher. *J. Am. Chem. Soc.* **129**, 3842–3843 (2007).
17. Biro, S. M. & Rebek, J. J. Structure and binding properties of water-soluble cavitands and capsules. *Chem. Soc. Rev.* **36**, 93–104 (2007).
18. Ballester, P. Anion binding in covalent and self-assembled molecular capsules. *Chem. Soc. Rev.* **39**, 3810–3830 (2010).
19. Hapiot, F., Tilloy, S. & Monflier, E. Cyclodextrins as supramolecular hosts for organometallic complexes. *Chem. Rev.* **106**, 767–781 (2006).
20. Adriaenssens, L. & Ballester, P. Hydrogen bonded supramolecular capsules with functionalized interiors: the controlled orientation of included guests. *Chem. Soc. Rev.* **42**, 3261–3277 (2010).
21. Dsouza, R. N., Pischel, U. & Nau, W. M. Fluorescent dyes and their supramolecular host/guest complexes with macrocycles in aqueous solution. *Chem. Rev.* **111**, 7941–7980 (2011).
22. Yagai, S. & Kitamura, A. Recent advances in photoresponsive supramolecular self-assemblies. *Chem. Soc. Rev.* **37**, 1520–1529 (2008).
23. Rananaware, A. *et al.* Photomodulation of fluoride ion binding through anion- $\pi$  interactions using a photoswitchable azobenzene system. *Sci. Rep.* **6**, 22928 (2016).
24. Natali, M. & Giordani, S. Molecular switches as photocontrollable “smart” receptors. *Chem. Soc. Rev.* **41**, 4010–4029 (2012).
25. Dube, H., Ajami, D. & Rebek, J. J. Photochemical control of reversible encapsulation. *Angew. Chem.* **49**, 3192–3195 (2010).
26. Clever, G. H., Tashiro, S. & Shionoya, M. Light-triggered crystallization of a molecular host-guest complex. *J. Am. Chem. Soc.* **132**, 9973–9975 (2010).
27. Han, M. *et al.* Light-triggered guest uptake and release by a photochromic coordination cage. *Angew. Chem. Int. Ed.* **52**, (1319–1323) (2013).
28. Kishi, N. *et al.* Facile catch and release of fullerenes using a photoresponsive molecular tube. *J. Am. Chem. Soc.* **135**, 12976–12979 (2013).
29. Yuan, K., Guo, Y.-J. & Zhao, X. A novel photo-responsive azobenzene-containing nanoring host for fullerene-guest facile encapsulation and release. *Phys. Chem. Chem. Phys.* **16**, 27053–27064 (2014).
30. Liang, X., Asanuma, H. & Komiyama, M. Photoregulation of DNA triplex formation by azobenzene. *J. Am. Chem. Soc.* **124**, 1877–1883 (2002).
31. Liang, X. *et al.* NMR study on the photoresponsive DNA tethering an azobenzene. Assignment of the absolute configuration of two diastereomers and structure determination of their duplexes in the trans-form. *J. Am. Chem. Soc.* **125**, 16408–16415 (2003).
32. Liu, M., Asanuma, H. & Komiyama, M. Azobenzene-tethered T7 promoter for efficient photoregulation of transcription. *J. Am. Chem. Soc.* **128**, 1009–1015 (2006).
33. Yan, Y., Wang, X., Chen, J. I. L. & Ginger, D. S. Photoisomerization quantum yield of azobenzene-modified DNA depends on local sequence. *J. Am. Chem. Soc.* **135**, 8382–8387 (2013).
34. Yuan, K., Guo, Y. J. & Zhao, X. A novel photo-responsive azobenzene-containing nanoring host for fullerene-guest facile encapsulation and release. *Phys. Chem. Chem. Phys.* **16**, 27053–27064 (2014).
35. Banerjee, I. A., Yu, L. & Matsui, H. Application of host-guest chemistry in nanotube-based device fabrication: photochemically controlled immobilization of azobenzene nanotubes on patterned r-CD monolayer/Au substrates via molecular recognition. *J. Am. Chem. Soc.* **125**, 9542–9543 (2003).
36. Shirai, Y. *et al.* Synthesis and photoisomerization of fullerene- and oligo(phenylene ethynylene)-azobenzene derivatives. *ACS Nano* **2**, 97–106 (2008).
37. Zhao, Y. L. & Stoddart, J. F. Azobenzene-based light-responsive hydrogel system. *Langmuir* **25**, 8442–8446 (2009).
38. Chen, S. L., Chu, C. C. & Hsiao, V. K. S. Reversible light-modulated photoluminescence from azobenzene-impregnated porous silicon. *J. Mater. Chem. C* **1**, 3529–3531 (2013).
39. Zhang, Y. *et al.* Enhancement of the photoresponse in organic field-effect transistors by incorporating thin DNA layers. *Angew. Chem. Int. Ed.* **53**, 244–249 (2014).
40. Commins, P. & Garcia-Garibay, M. A. Photochromic molecular gyroscope with solid state rotational states determined by an azobenzene bridge. *J. Org. Chem.* **79**, 1611–1619 (2014).
41. Li, Z., Xue, W., Liu, G., Liu, S. H. & Yin, J. Switchable azo-macrocycle: from molecules to functionalisation. *Supramolecular Chemistry* **26**, 54–65 (2014).
42. Zhang, X., Zhao, H., Tian, D., Deng, H. & Li, H. A Photoresponsive wettability switch based on a dimethylamino calix[4]arene. *Chem. Eur. J.* **20**, 9367–9371 (2014).
43. Kienzler, M. A. *et al.* A Red-shifted, fast-relaxing azobenzene photoswitch for visible light control of an ionotropic glutamate receptor. *J. Am. Chem. Soc.* **135**, 17683–17686 (2013).
44. Reiter, A., Skerra, A., Trauner, D. & Schiefner, A. A photoswitchable neurotransmitter analogue bound to its receptor. *Biochemistry* **52**, 8972–8974 (2013).
45. Frank, J. A. *et al.* Photoswitchable fatty acids enable optical control of TRPV1. *Nat. Commun.* **6**, 7188 (2015).
46. Yuan, K., Dang, J.-S., Guo, Y.-J. & Zhao, X. Theoretical prediction of the host-guest interactions between novel photoresponsive nanorings and C<sub>60</sub>: A strategy for facile encapsulation and release of fullerene. *J. Comput. Chem.* **36**, 518–528 (2015).
47. Yamamura, M., Saito, T. & Nabeshima, T. phosphorus-containing chiral molecule for fullerene recognition based on concave/convex interaction. *J. Am. Chem. Soc.* **136**, 14299–14306 (2014).
48. Litvinov, A. L. *et al.* Molecular complexes of fullerene C<sub>60</sub> with aromatic hydrocarbons containing flexible phenyl substituents. *CrystEngComm* **4**, 618–622 (2002).
49. Anuradha, L. A., Suryawanshi, D. D., Al, K. M. & Bhosale, S. V. Right handed chiral superstructures from achiral molecules: self-assembly with a twist. *Sci. Rep.* **5**, 15652 (2015).
50. Mei, J., Leung, N. L. C., Kwok, R. T. K., Lam, J. W. Y. & Tang, B. Z. Aggregation-Induced Emission: Together We Shine, United We Soar! *Chem. Rev.* **115**, 11718–11940 (2015).
51. Catti, L., Kishida, N., Kai, T., Akita, M. & Yoshizawa, M. When this manuscript was under review the following work on the similar topic was published. Polyaromatic nanocapsules as photoresponsive hosts in water. *Nat. Commun.* **10**, 1948 (2019).

## Acknowledgements

S.V.B. (GU) acknowledges Financial Support and Professorship from the UGC-FRP. S.B. acknowledges DST-SERB for a research grant (EMR/2017/003720). M.S. is funded by a UGC SRF fellowship, SAR by a doctoral fellowship from IISER Kolkata and M Saha by a DST INSPIRE fellowship. D.N. and R.W.J. acknowledges UGC for Junior Research Fellowship's.

### Author Contributions

M.S. performed characterisation of host-guest complexation by mean of UV-vis, fluorescence and NMR spectroscopy. D.N.N., A.R. and R.W.J. performed syntheses and structure spectroscopic characterisation of the compounds used in this study. S.A.R. and M.S. performed crystallographic studies and the computational analysis. S.B. and S.V.B. plan directed the research and interpreted and analyse the data and drafted the manuscript. All co-authors reviewed the manuscript.

### Additional Information

**Supplementary information** accompanies this paper at <https://doi.org/10.1038/s41598-019-46242-4>.

**Competing Interests:** The authors declare no competing interests.

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019