

# Arbuscular Mycorrhizal (AM) fungi for sustainable agriculture

**K. M. RODRIGUES AND B. F. RODRIGUES**



*J. Mycopathol. Res.* 58(3) : 135-140, 2020;  
ISSN 0971-3719

© Indian Mycological Society,  
Department of Botany,  
University of Calcutta,  
Kolkata 700 019, India

***This article is protected by copyright and all other rights under the jurisdiction of the Indian Mycological Society. The copy is provided to the author(s) for internal non-commercial research and educational purposes.***

---

## REVIEW

# Arbuscular Mycorrhizal (AM) fungi for sustainable agriculture

---

**K. M. RODRIGUES AND B. F. RODRIGUES\***

*Department of Botany, Goa University, Goa 403 206, India.*

---

Received : 19.06.2020

Accepted : 30.07.2020

Published : 26.10.2020

---

Arbuscular mycorrhizal (AM) fungi are obligate biotrophic symbionts that form a mutualistic association with plant roots. They are directly involved in plant mineral nutrition and environmental stress tolerance. Most crop plants are symbiotic with AM fungi, and the ability of AM fungal symbiosis to improve crop production is widely recognized. This paper highlights the efficacy of AM fungi as a vital component of sustainable crop production systems, and its prospective for exploitation as an on-farm agro-input.

**Key words :** Sustainable agriculture, arbuscular mycorrhizal fungi, ecological processes, AM fungal diversity, biofertilizer

---

## INTRODUCTION

Sustainable agriculture is a method of agricultural production, which chiefly excludes the usage of synthetic agro-inputs (chemical fertilizers and pesticides) and to the maximum extent relies upon the application of environmentally friendly organic agro-inputs (organic wastes and beneficial microbes), thereby promoting and enhancing overall agroecosystem health including biodiversity, biological cycles, and soil biological activity. Some of the latest estimates predict that by 2050, the world's population will reach 9.1 billion, 34 percent higher than today, hence emphasizing the need to increase agricultural productivity (FAO, 2009). The global demand for crops certainly accentuates the necessity to implement eco-friendly agrarian management practices for sustainable agricultural production (Islam and Katoh, 2017).

As a result, the scientific community all over the world is continuously searching for economically viable, socially safe, and environmentally sustainable on-farm resources. Thus the focus shifts increasingly towards the role of soil microbial diversity in general and particularly arbuscular

mycorrhizal (AM) fungi in achieving this in a sustainable manner (Bender *et al.*, 2016; Thirkell *et al.*, 2017; Sosa-Hernández *et al.*, 2019). Soil microbes play an imperative role in most biological transformations, including nutrient recycling, which facilitates the subsequent establishment of plants. Fungi are the most efficient soil microorganisms involved in soil-structure stabilization, and AM fungi often comprise the significant fraction of the soil microbiome.

### **Arbuscular Mycorrhizal (AM) fungi**

Arbuscular mycorrhiza (AM) which constitutes 72% of various types of mycorrhizas (van der Heijden *et al.*, 2015; Brundrett, 2017), is ecologically an important type of mycorrhiza which is most widely commercially exploited in agricultural production systems (Owen *et al.*, 2015; Rillig *et al.*, 2016). AM fungi are ubiquitous soil fungi that form a mutualistic relationship with most land plants. These fungi are a monophyletic lineage of obligate mycobionts belonging to phylum Glomeromycota. As the phylum is evolutionarily an ancient form of symbiosis in plants, about 90% of existing plant species are mycorrhizal, including many crops (Moëne-Loccoz *et al.*, 2015; Brundrett and Tedersoo, 2018), therefore leading to an array of positive effects in both natural ecosystem and agricultural biotas (van der Heijden *et al.*, 2015).

---

\*Corresponding author : felinov@gmail.com

### ***AM fungal-plant root interaction for the establishment of symbiosis***

AM fungi are present as chlamydospores (spores) colonizing plant roots in the rhizosphere. In the asymbiotic fungal growth phase, a chlamydospore will germinate by forming aseptate and thick-walled hypha (permanent hypha). Thin-walled short-lived hyphae arise from the permanent hypha and grow along with the roots. The exploratory hyphal growth changes dramatically in the presence of signals derived from plant root exudates (strigolactones) resulting in the development of pre-symbiotic fungal growth phase characterized by profuse hyphal branching, increased physiological activity and continued hyphal growth (López-Ráez *et al.*, 2011). It takes one to several weeks for the establishment of host root contact by the fungal hyphae. Once the fungus-plant root contact is established, changes in the fungal morphology and metabolism occur radically, marking the initiation of the symbiotic phase. AM fungi produce mycorrhiza (Myc) factors that can induce calcium oscillations in root epidermal cells and activate plant AM symbiosis-related genes (Bonfante and Genre, 2010). At the point of contact with the plant root, the hyphae bear appressoria and penetrate the root hair or epidermis to reach the cortex, bifurcating intra- and inter-cellularly from the point of entry, leading to the development of intra-radical structures (hyphae, arbuscules, vesicles) in the cortical cells of the root and extra-radical structures (hyphae and spores) in the soil. Arbuscules are repeated dichotomously branched tree-like structures that are formed intra-cellularly within the cortical cells. They act as a site of nutrient exchange and are responsible for transferring carbohydrates from the host plant tissues to the fungus, and supplying mineral nutrients and water from the fungus to the host plant (Rich *et al.*, 2017). Vesicles are thin or thick-walled oval structures that are formed inter- and intra-cellularly within the cortical cells. They act as storage structures, storing materials like lipids, poly-phosphate, and other minerals. Vesicles also function as propagules (chlamydospores) in older roots. AM fungi also produce a mycelial network outside the root in the soil known as the extra-radical mycelium, where spores and sporocarps are produced along with mycelium that serves as propagules to complete the AM fungal life cycle (Barea *et al.*, 2013; Gupta, 2017). The AM fungal inoculum, which includes soil-based propagules

(spores, extra-radical hyphae, and colonized root fragments) allows AM fungi to colonize new plant roots through an established extra-radical mycelial network in the soil (Muller *et al.*, 2017).

### **Agro-ecological functions of AM fungi**

AM fungal association is characterized by a bidirectional nutrient flux wherein the mycobiont helps the phytobiont in the acquirement of soil nutrients (mainly P) while the phytobiont provides photosynthates (carbon sources) to the mycobiont (Wipf *et al.*, 2019). The basis of this symbiotic association is the ability of AM fungi to form fine hyphae with a more favourable surface area-to-volume ratio for nutrient uptake and to secrete enzymes or organic acids to mobilize nutrients (Owen *et al.*, 2015). AM fungi are primarily known for their capability to increase plant growth and productivity by augmenting macro- and micro-nutrient acquisition (Smith and Smith, 2011; Hart and Forsythe 2012; Pellegrino and Bedini 2014). AM fungal symbiosis allows better plant phosphorus uptake from soils with low levels of available P (van der Heijden *et al.*, 2015); they help in the uptake of nitrogen from the soil and increase the utilization of different forms of N (Jacott *et al.*, 2017); they also contribute in the uptake of calcium, iron, manganese, zinc, copper, sulphur, potassium, magnesium and aluminium in host plants (Hart *et al.*, 2017). AM fungi are most beneficial in improving plant nutrient acquisition, especially in low-fertility soils. It is widely believed that AM fungi can serve as an alternative for inorganic fertilizers because their application can effectively reduce the quantitative use of chemical fertilizer input, especially P (Ortas, 2012).

Besides nutrient acquisition, AM fungi also influence a wide range of other host benefits and ecosystem processes. Such as resistance to several abiotic stress factors (Sun *et al.*, 2018); safeguarding the host plants from pathogenic infections (Jung *et al.*, 2012), hence AM fungi can be utilized as biocontrol agents. AM fungal symbiosis also stimulates the synthesis of several plant secondary metabolites *viz.*, polyphenols, phytosterols, vitamins, lignins, terpenoids, and carotenoids for increased plant tolerance to abiotic and biotic stresses. The formation of phytotoxins, chitinases, glucanases, peroxidases, PR proteins, phenolics, and hydroxyl proline-rich glycoproteins as defense responses also increases in host plants

upon association with AM fungi (Smith *et al.*, 2010). Initiation of AM fungal symbiosis causes the host plants to secrete organic acids, polysaccharides, phenolic compounds, amino acids, and enzymes like phosphatases as exudates from the root system, which favours the growth of AM fungi and host plant. This also has a positive effect on the associated microbiome functioning (Gupta *et al.*, 2018a, b). Improved soil stability, binding, and water retention by the AM fungal hyphae through the ramifying extra-radical mycelial network and secretion of hydrophobic “sticky” proteinaceous substance (glomalin) in the soil, consequently reducing soil erosion. AM fungi, through their extensive extra-radical mycelial network, interconnect several unrelated individual plant species, consequently impacting the function and biodiversity of the entire ecosystem (Bonfante and Genre, 2010). AM fungi also positively influence nutrient cycling, energy flow, and plant establishment in disturbed ecosystems.

Moreover, AM fungi are involved in bioremediation of soil. They help in alleviating heavy metal toxicity to plants by reducing metal translocation from root to shoot. Therefore, they contribute to revegetation and restoration of disturbed or contaminated lands. Thus, AM fungi are imperative endosymbionts playing an efficient role in plant growth and productivity along with the functioning of the ecosystems. They are of immense significance for sustainable crop improvement (Gianinazzi *et al.*, 2010).

### **Significance of AM fungal diversity and dominance in sustainable agriculture**

Ecosystems and plant life are widely influenced by the functionally diverse nature of AM fungi, the symbiosis is considered as a tripartite relationship between plant, fungus, and soil where the fungus creates a close link between the soil and the plant. AM fungi co-exist as assemblages of mixed species in terrestrial ecosystems with certain species being dominant, and a plant may be colonized by several AM species at one time. Most of the AM fungal species are widespread occurring in different terrestrial habitats and are considered ‘generalists’, however some species appear to be restricted to particular ecosystem types and are considered ‘specialists’. According to the latest estimates, approximately 300 species of AM fungi have been reported from all over the world (Öpik

*et al.*, 2013). In India, more than 161 species have been reported (Gupta *et al.*, 2017). Studies from India showed that *Funneliformis mosseae* is the most widely distributed species, and the genus *Glomus* is represented by the highest number of species (Gupta *et al.*, 2014).

Different soil types showing variation in AM species can be characterized by AM fungal community structure (Oehl *et al.*, 2010). AM taxa differ significantly in their life histories and are believed to be nonspecific about their capability to infect and colonize different plant species, although there may be exceptions. There is also variation in response to soil structure, mineral acquisition, plant health, growth rate, biomass allocation, and symbiotic effects. AM species diversity is more distinct in undisturbed ecosystems compared to disturbed ecosystems as in undisturbed ecosystems. There is a greater degree of variability in terms of critical determinants. AM diversity and abundance can be affected by various factors/determinants such as habitat type, edaphic conditions, climatic or seasonal variations, host genotype, and vegetation cover. AM dominance can also be affected by the severity and extent of disturbance in a habitat (Bhatia *et al.*, 2013). Plant diversity and productivity are enhanced by AM symbiosis and AM species richness. An increase in AM fungal diversity results in an increase in species richness and hence higher plant productivity. This suggests that changes in below-ground AM fungal diversity can affect changes in above-ground plant diversity and productivity. Assessment of AM fungal diversity is essential if the benefits associated with the symbiosis are to be exploited. Knowledge of AM species diversity in functioning ecosystems is crucial for the development of inocula for agricultural and horticultural crops, and revegetation of degraded ecosystems.

### **AM fungi as a biofertilizer for sustainable crop production systems**

AM fungi form a symbiotic association with most agricultural crops *viz.*, *Triticum* (wheat), *Glycine max* (soybean), *Phaseolus vulgaris* (beans), *Solanum lycopersicum* (tomato), *Sorghum bicolor* (sorghum), *Capsicum annum* (red peppers) and *Daucus carota* (carrot) (Öpik *et al.*, 2010). As a result, they have gained growing attention as ecosystem engineers and bio-inoculants (Fitter *et*

*al.*, 2011). Currently, AM fungal bio-inoculants are being increasingly considered in agriculture, horticulture, and forestry, as well as for environmental reclamation and remediation, to increase overall crop health and yield, and to limit the usage of agrochemicals. Many fields and greenhouse studies have been limited to the single inoculation of one of the following three species *viz.*, *Rhizoglyphus intraradices*, *R. irregulare*, and *Funneliformis mosseae* (Krüger *et al.*, 2012; Pellegrino and Bedini, 2014; Berruti *et al.*, 2015). These AM species are very generalist symbionts that colonize a diversity of host plants. They can be quickly and massively propagated, survive long-term storage, and have a wide geographical distribution all over the world (Öpik *et al.*, 2010). Several reports have also highlighted that different isolates within the same species, rather than different species, can cause larger variations in host plant response (Angelard *et al.*, 2010). The abovementioned characteristics have made these AM fungal species suitable as premium inoculum components and their application by farmers is increasing (Gianinazzi, 2014; Berruti *et al.*, 2015; Hijri, 2016).

AM fungal inoculum is commercially available in a variety of forms ranging from high concentrations of AM fungal propagules in carrier materials to potting media containing inoculum at low concentrations (Douds *et al.*, 2010). Interest in AM fungi has focused on finding a viable method to optimize the production of high quality AM fungal inoculum to use as a bio-inoculant in cropping systems (Ijdo *et al.*, 2011). *In vitro* culture of AM fungi (monoxenic cultivation) is now a reality offering a several-fold increase in mass production of effective propagules over conventional bulking techniques. This system of AM inoculum production provides potentially high and economically attractive options to chemical fertilization, thus emphasizing its potential importance in sustainable agriculture.

AM fungal symbiosis contributes to sustainable agriculture principally by serving as an advantageous nutrient sink in which AM inoculated or mycorrhizal plants have increased growth and yield as compared to un-inoculated or non-mycorrhizal plants. This is due to better nutrient extraction and uptake capacity and improved soil properties. Thus, AM fungi impart beneficial effects to crops in terms of growth and productivity by

increasing favourable microbial interactions and resistance to unfavourable biotic or abiotic conditions (Verbruggen *et al.*, 2010; Lenoir *et al.*, 2016).

## CONCLUSION

AM fungi are universal, complex, ancient microorganisms constituting a significant component not only of plant life but also of agro-ecological production. It is clear they significantly impact every terrestrial niche across almost every biome influencing plant functioning. The ecological services provided by AM fungi and the extent of those services at the interface between biotic and abiotic elements suggest that AM fungi are primary ecosystem facilitators “keystone mutualists”.

In sustainable crop production, to obtain maximum beneficial effects from AM fungi emphasis must be on green agricultural practices or organic farming system relying on continuous rotational crops, cover crops, reduced tillage, application of organic fertilizers and manures. This would support the occurrence and functioning of AM fungal diversity as well as nutrient cycling in the long term thereby sustaining higher crop productivity without compromising the soil health.

## REFERENCES

- Angelard, C., Colard, A., Niculita-Hirzel, H., Croll, D. and Sanders, I.R. 2010. Segregation in a mycorrhizal fungus alters rice growth and symbiosis-specific gene transcription. *Curr. Biol.* **20**: 1216-1221.
- Barea, J.M., Pozo, M.J., Lopez-Raez, J.A., Aroca, R., Ruiz-Lozano, J.M., Ferrol, N., Azcón, R. and Azcón-Aguilar, C. 2013. Arbuscular mycorrhizas and their significance in promoting soil-plant systems sustainability against environmental stresses. Rodelas, B. and González-López, J. (Eds.) In: *Beneficial plant-microbial interactions: ecology and applications*. pp: 353-387. CRC Press, USA.
- Bender, S.F., Wagg, C. and van der Heijden, M.G.A. 2016. An underground revolution: biodiversity and soil ecological engineering for agricultural sustainability. *Trends Ecol. Evol.* **31** (6): 440-452.
- Berruti, A., Lumini, E., Balestrini, R. and Bianciotto, V. 2015. Arbuscular mycorrhizal fungi as natural biofertilizers: let's benefit from past successes. *Front. Microbiol.* **6**: 1559.
- Bhatia, N.P., Sundari, K. and Adholeya, A. 2013. Diversity and selective dominance of vesicular-arbuscular mycorrhizal fungi. Mukerji, K.G. (Ed.) In: *Concepts in mycorrhizal research*. pp: 133-178. Springer Science and business media.
- Bonfante, P. and Genre, A. 2010. Mechanisms underlying beneficial plant-fungus interactions in mycorrhizal symbiosis. *Nat. Commun.* **1**: 48.
- Brundrett, M.C. 2017. Global diversity and importance of mycorrhizal and non-mycorrhizal plants. Tedersoo, L. (Ed.) In: *Biogeography of mycorrhizal symbiosis*. pp: 533-556. Springer, Cham.

- Brundrett, M.C. and Tedersoo, L. 2018. Evolutionary history of mycorrhizal symbioses and global host plant diversity. *New Phytol.* **220**: 1108-1115.
- Douds, D.D. Jr., Nagahashi, G. and Hepperly, P.R. 2010. On-farm production of inoculum of indigenous arbuscular mycorrhizal fungi and assessment of diluents of compost for inoculum production. *Bioresour. Technol.* **101**: 2326-2330.
- FAO. 2009. Report of the FAO Expert Meeting on How to Feed the World in 2050. FAO, Rome.
- Fitter, A.H., Helgason, T. and Hodge, A. 2011. Nutritional exchanges in the arbuscular mycorrhizal symbiosis: implications for sustainable agriculture. *Fungal Biol. Rev.* **25**: 68-72.
- Gianinazzi, S., Golotte, A., Binet, M. N., Van Tuinen, D., Redecker, D. and Wipf, D. 2010. Agroecology: the key role of arbuscular mycorrhizas in ecosystem services. *Mycorrhiza* **20**: 519-530.
- Gianinazzi, S.G. 2014. Domestication of beneficial soil microorganisms: an innovative technology for agriculture. *Int. Congr. Mycorrhizae, Marrakesh.* p. 26.
- Gupta, M. 2017. Differential response of arbuscular mycorrhizal sporocarps in long-term trap culturing. *Phytomorphology* **67**: 1-11.
- Gupta, M.M., Aggarwal, A. and Asha 2018a. From mycorrhizosphere to rhizosphere microbiome: the paradigm shift. Giri, B., Prasad, R. and Varma, A. (Eds.) In: *Root biology. Soil biology.* Springer, Cham. pp: 487-500.
- Gupta, M.M., Gupta, A. and Kumar, P. 2018b. Urbanization and biodiversity of arbuscular mycorrhizal fungi: the case study of Delhi, India. *Rev. Biol. Trop.* **66**: 1563-1574.
- Gupta, M.M., Naqvi, N. and Kumar, P. 2017. AMF-centralized database of arbuscular mycorrhizal distribution, phylogeny and taxonomy. *J. Recent Adv. Appl. Sci.* **30**: 1-5.
- Gupta, M.M., Naqvi, N.S. and Singh, V.K. 2014. The state of arbuscular mycorrhizal fungal diversity in India: an analysis. *Sydowia* **66**: 265-288.
- Hart, M.M. and Forsythe, J.A. 2012. Using arbuscular mycorrhizal fungi to improve the nutrient quality of crops; nutritional benefits in addition to phosphorus. *Sci. Hortic.* **148**: 206-214.
- Hart, M.M., Antunes, P.M. and Abbott, L.K. 2017. Unknown risks to soil biodiversity from commercial fungal inoculants. *Nat. Ecol. Evol.* **1**: 115.
- Hijri, M. 2016. Analysis of a large dataset of mycorrhiza inoculation field trials on potato shows highly significant increases in yield. *Mycorrhiza* **26**: 209-214.
- Ijdo, M., Cranenbrouck, S. and Declerck, S. 2011. Methods for large-scale production of AM fungi: past, present, and future. *Mycorrhiza* **21**: 1-16.
- Islam, M.Z. and Katoh, S. 2017. The effect of arbuscular mycorrhizal fungi and *Gliocladium* fungi on the yield of small green pepper (*Capsicum annuum*) grown by sustainable agriculture. *Agricultural Sciences*, **8**: 1296-1314.
- Jacott, C.N., Murray, J.D. and Ridout, C.J. 2017. Trade-offs in arbuscular mycorrhizal symbiosis: disease resistance, growth responses and perspectives for crop breeding. *Agronomy* **7** (4): 75.
- Jung, S.C., Martinez-Medina, A., Lopez-Raez, J.A. and Pozo, M.J. 2012. Mycorrhiza-induced resistance and priming of plant defenses. *J. Chem. Ecol.* **38**: 651-664.
- Krüger, M., Krüger, C., Walker, C., Stockinger, H. and Schüßler, A. 2012. Phylogenetic reference data for systematics and phylotaxonomy of arbuscular mycorrhizal fungi from phylum to species level. *New Phytol.* **193**: 970-984.
- Lenoir, I., Lounes-Hadj Sahaoui, A. and Fontaine, J. 2016. Arbuscular mycorrhizal fungal-assisted phytoremediation of soil contaminated with persistent organic pollutants: a review. *Eur. J. Soil Sci.* **67**: 624-640.
- López-Ráez, J.A., Charnikhova, T., Fernández, I., Bouwmeester, H. and Pozo, M.J. 2011. Arbuscular mycorrhizal symbiosis decreases strigolactone production in tomato. *J. Plant Physiol.* **168**: 294-297.
- Moënné-Loccoz, Y., Mavingui, P., Combes, C., Normand, P. and Steinberg, C. 2015. Microorganisms and biotic interactions. Bertrand, J.C., Caumette, P., Lebaron, P., Matheron, R., Normand, P. and Sime-Ngando, T. (Eds.) In: *Environmental microbiology: fundamentals and applications.* Springer, Dordrecht. pp: 395-444.
- Muller, A., Ngwene, B., Peiter, E. and George, E. 2017. Quantity and distribution of arbuscular mycorrhizal fungal storage organs within dead roots. *Mycorrhiza* **27**: 201-210.
- Oehl, F., Laczko, E., Bogenrieder, A., Stahr, K., Bösch, R., van der Heijden, M.G.A. and Sieverding, E. 2010. Soil type and land use intensity determine the composition of arbuscular mycorrhizal fungal communities. *Soil Biol. Biochem.* **42**: 724-738.
- Öpik, M., Davison, J., Moora, M. and Zobel, M. 2013. DNA-based detection and identification of Glomeromycota: the virtual taxonomy of environmental sequences. *Botany* **92**: 135-147.
- Öpik, M., Vanatoa, A., Vanatoa, E., Moora, M., Davison, J., Kalwij, J.M. and Zobel, M. 2010. The online database MaarjAM reveals global and ecosystemic distribution patterns in arbuscular mycorrhizal fungi (Glomeromycota). *New Phytol.* **188**: 223-241.
- Ortas, I. 2012. The effect of mycorrhizal fungal inoculation on plant yield, nutrient uptake and inoculation effectiveness under long-term field conditions. *Field Crops Res.* **125**: 35-48.
- Owen, D., Williams, A.P., Griffith, G.W. and Withers, P.J.A. 2015. Use of commercial bio-inoculants to increase agricultural production through improved phosphorus acquisition. *Appl. Soil Ecol.* **86**: 41-54.
- Pellegrino, E. and Bedini, S. 2014. Enhancing ecosystem services in sustainable agriculture: biofertilization and biofortification of chickpea (*Cicer arietinum* L.) by arbuscular mycorrhizal fungi. *Soil Biol. Biochem.* **68**: 429-439.
- Rich, M.K., Nouri, E., Courty, P.E. and Reinhardt, D. 2017. Diet of arbuscular mycorrhizal fungi: bread and butter? *Trends Plant Sci.* **22** (8): 652-660.
- Rillig, M.C., Sosa-Hernández, M.A., Roy, J., Aguilar-Trigueros, C.A., Vályi, K. and Lehmann, A. 2016. Towards an integrated mycorrhizal technology: harnessing mycorrhiza for sustainable intensification in agriculture. *Front Plant Sci.* **7**: 1625.
- Smith, S.E. and Smith, F.A. 2011. Roles of arbuscular mycorrhizas in plant nutrition and growth: new paradigms from cellular to ecosystem scales. *Annu. Rev. Plant Biol.* **62**: 227-250.
- Smith, S.E., Facelli, E., Pope, S. and Smith, F.A. 2010. Plant performance in stressful environments: interpreting new and established knowledge of the roles of arbuscular mycorrhizas. *Plant Soil.* **326** (1-2): 3-20.
- Sosa-Hernández, M.A., Leifheit, E.F., Ingrassia, R. and Rillig, M.C. 2019. Subsoil arbuscular mycorrhizal fungi for sustainability and climate-smart agriculture: a solution right under our feet? *Front. Microbiol.* **10**: 744.
- Sun, Z., Song, J., Xin, X., Xie, X. and Zhao, B. 2018. Arbuscular mycorrhizal fungal proteins 14-3-3- are involved in arbuscule formation and responses to abiotic stresses during AM symbiosis. *Front. Microbiol.* **5**: 9-19.
- Thirkell, T.J., Charters, M.D., Elliott, A.J., Sait, S.M. and Field, K.J. 2017. Are mycorrhizal fungi our sustainable saviours? Considerations for achieving food security. *J. Ecol.* **105**: 921-929.
- van der Heijden, M.G.A., Martin, F.M., Selosse, M.A. and Sanders, I.R. 2015. Mycorrhizal ecology and evolution: the past, the present, and the future. *New Phytol.* **205**: 1406-1423.

- Verbruggen, E., Røling, W.F.M., Gamper, H.A., Kowalchuk, G.A., Verhoef, H.A. and van der Heijden, M.G.A. 2010. Positive effects of organic farming on below-ground mutualists: large-scale comparison of mycorrhizal fungal communities in agricultural soils. *New Phytol.* **186**: 968-979.
- Wipf, D., Krajinski, F., van Tuinen, D., Recorbet, G. and Courty, P.E. 2019. Trading on the arbuscular mycorrhiza market: from arbuscules to common mycorrhizal networks. *New Phytol.* **223**: 1127-1142.